

On periodic solutions of second-order differential equations with attractive–repulsive singularities

Robert Hakl¹ and Pedro J. Torres²

⁽¹⁾ *Institute of Mathematics AS CR, Žitkova 22, 616 62 Brno, Czech Republic*

⁽²⁾ *Departamento de Matemática Aplicada, Facultad de Ciencias, Universidad de Granada, Campus de Fuentenueva s/n, 18071 Granada, Spain.*

Abstract

Sufficient conditions for the existence of a solution to the problem

$$\begin{aligned} u''(t) &= \frac{g(t)}{u^\mu(t)} - \frac{h(t)}{u^\lambda(t)} + f(t) && \text{for a. e. } t \in [0, \omega], \\ u(0) &= u(\omega), && u'(0) = u'(\omega) \end{aligned}$$

are established.

2000 Mathematics Subject Classification: 34B18, 34C25

Key words and phrases: Second-order ordinary differential equation, singular equation, periodic solution, positive solution.

Introduction

In this paper, we are concerned with the periodic problem

$$u''(t) = \frac{g(t)}{u^\mu(t)} - \frac{h(t)}{u^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega], \quad (0.1)$$

$$u(0) = u(\omega), \quad u'(0) = u'(\omega), \quad (0.2)$$

where $g, h \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$, $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, and $\lambda, \mu > 0$. By a solution to (0.1), (0.2) we understand a function $u \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ satisfying (0.1). Special cases of the

¹Supported by the Academy of Sciences of the Czech Republic, Institutional Research Plan No. AV0Z10190503

²Supported by Ministerio de Educación y Ciencia, Spain, project MTM2008-02502, and by Junta de Andalucía, Spain, Project FQM2216.

equation (0.1) are

$$u''(t) = \frac{g(t)}{u^\mu(t)} - \frac{h(t)}{u^\lambda(t)} \quad \text{for a. e. } t \in [0, \omega], \quad (0.3)$$

$$u''(t) = -\frac{h(t)}{u^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega], \quad (0.4)$$

$$u''(t) = \frac{g(t)}{u^\mu(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (0.5)$$

In the related literature, it is said that (0.4) has an attractive singularity, whereas (0.5) has a repulsive singularity. The interest on this type of equations began with the paper of Lazer and Solimini [7], where the authors provide necessary and sufficient conditions for existence of periodic solutions of eq. (0.4) and (0.5) with constant positive functions h , g and a continuous forcing term f . Their proofs can be easily extended to the case when the function h , resp. g is bounded from below by some positive constant (see the generalized results presented in the paper of Habets and Sanchez [3]), but in their arguments this hypothesis is essential and cannot be omitted. In the repulsive case, a strong force assumption ($\mu \geq 1$) is also essential.

The equation (0.3) is interesting due to a mixed type of singularity on the right-hand side. Since the functions g and h are possibly zero on some sets of positive measure, the singularity may combine attractive and repulsive effects. If h, g are positive constants, the singular term can be regarded as a generalized Lennard-Jones force or van der Waals attraction/repulsion force and it is widely use in Molecular Dynamics to model the interaction between atomic particles (see for instance [4, 9, 12, 15] and the references therein). In a different physical context, a periodic solution of equation (0.3) is equivalent to a matter-wave breather in a Bose-Einstein condensate with a periodic control of the scattering length (the mathematical model is a nonlinear Schrödinger equation with a cubic term, then the method of moments leads to the study of a particular case of (0.3), see [8] for more details. Finally, a third different range of applicability is the evolution of optical pulses in dispersion-managed fiber communication devices [6].

In spite of the variety of physical applications, the analysis of differential equations with mixed singularities is at this moment very incomplete, and few references can be cited (see [1, 5, 13]) if compared with the large number of references devoted to singular equations, either of attractive or repulsive type (see the review [10] and the references therein). Our main purpose in this paper is to contribute to the literature trying to fill partially this gap in the study of singularities of mixed type with an approach that should be useful as a starting point for further studies. Incidentally, our main results can be applied to the original Lazer-Solimini equations both in the attractive and in the repulsive case, giving new sufficient conditions for existence of periodic solutions when the functions h and g are possibly zero on the sets of positive measure.

The structure of the paper is as follows: Section 2 contains the tools needed in the proofs. In Section 3 we state and prove the main results and develop some corollaries for the equation with a singularity of mixed type. To illustrate the results, an application to

the dynamics of a trapless Bose-Einstein condensate is given. This model and related ones deserve a different treatment more oriented to a physical audience, that will be performed elsewhere. Finally, due its relevance in the related literature we have decided to devote Sections 4 and 5 to perform a comparative study of the equation with attractive (resp. repulsive) singularity. Along the paper, some open problems are posed. We feel that their consideration will bring light to this subject in the future.

The following notation is used throughout the paper:

\mathbb{R} is a set of all real numbers, $\mathbb{R}_+ = [0, +\infty[$, $[x]_+ = \max\{x, 0\}$, $[x]_- = \max\{-x, 0\}$.

$L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ is the Banach space of ω -periodic Lebesgue integrable functions $p : \mathbb{R}/\omega\mathbb{Z} \rightarrow \mathbb{R}$.

$AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ is a set of all ω -periodic functions $u : \mathbb{R}/\omega\mathbb{Z} \rightarrow \mathbb{R}$ such that u and u' are absolutely continuous.

$L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+) = \{p \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}) : p(t) \geq 0 \text{ for a. e. } t \in [0, \omega]\}$.

Notation 0.1. For the sake of brevity we will use the following notation throughout the paper:

$$\begin{aligned} G &= \int_0^\omega g(s) ds, & H &= \int_0^\omega h(s) ds, \\ F &= \int_0^\omega f(s) ds, & F_+ &= \int_0^\omega [f(s)]_+ ds, & F_- &= \int_0^\omega [f(s)]_- ds. \end{aligned}$$

Note that $F = F_+ - F_-$.

1 Auxiliary results

The proofs of our results rely on the method of upper and lower functions. The following two lemmas are classical and can be found, e.g., in [2, 14]. We introduce them in a form suitable for us.

Lemma 1.1. *Let there exist positive functions $\alpha, \beta \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ such that*

$$\alpha''(t) \geq \frac{g(t)}{\alpha^\mu(t)} - \frac{h(t)}{\alpha^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega], \quad (1.1)$$

$$\beta''(t) \leq \frac{g(t)}{\beta^\mu(t)} - \frac{h(t)}{\beta^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega], \quad (1.2)$$

$$\alpha(t) \leq \beta(t) \quad \text{for } t \in [0, \omega].$$

Then there exists at least one positive solution to (0.1), (0.2).

A function $\alpha \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ (resp. $\beta \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$) verifying (1.1) (resp. (1.2)) is called lower (resp. upper) function. When the order between the lower and the upper function is the inverse, an additional hypothesis is needed.

Definition 1.1. A function $\varphi \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$ is said to verify the property (P) if the implication

$$\left. \begin{array}{l} u \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}) \\ u''(t) + \varphi(t)u(t) \geq 0 \quad \text{for a. e. } t \in [0, \omega] \end{array} \right\} \implies u(t) \geq 0 \quad \text{for } t \in [0, \omega]$$

holds.

Lemma 1.2. Let there exist positive functions $\alpha, \beta \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ satisfying (1.1), (1.2) and

$$\beta(t) \leq \alpha(t) \quad \text{for } t \in [0, \omega].$$

Let, moreover, there exists $\varphi \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$ with the property (P) and such that

$$\frac{g(t)}{u^\mu(t)} - \frac{h(t)}{u^\lambda(t)} - \left(\frac{g(t)}{v^\mu(t)} - \frac{h(t)}{v^\lambda(t)} \right) \leq \varphi(t)(v(t) - u(t)) \quad \text{for a. e. } t \in [0, \omega], \quad (1.3)$$

whenever $\beta(t) \leq u(t) \leq v(t) \leq \alpha(t)$ for $t \in [0, \omega]$. Then there exists at least one positive solution to (0.1), (0.2).

Property (P) is just a maximum principle for the linear operator $Lu := u'' + \varphi(t)u$ with periodic boundary conditions, and it is equivalent to have a nonnegative Green function. The reference [14] provides sufficient conditions in the L^p -norm for $\varphi(t)$ to verify property (P). In particular, we have the following lemma.

Lemma 1.3. Let us assume that $\varphi \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$, $\varphi \not\equiv 0$, and at least one of the following conditions holds:

$$i) \quad \varphi(t) \leq \left(\frac{\pi}{\omega}\right)^2 \quad \text{for a. e. } t \in [0, \omega],$$

$$ii) \quad \int_0^\omega \varphi(t)dt \leq \frac{4}{\omega}.$$

Then, φ verifies the property (P).

To finish this section, we show a technical bound on the amplitude of oscillation of a periodic function.

Lemma 1.4. Given $v \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, then

$$M_v - m_v \leq \frac{\omega}{4} \int_0^\omega [v''(s)]_+ ds, \quad (1.4)$$

where

$$M_v = \max \{v(t) : t \in [0, \omega]\}, \quad m_v = \min \{v(t) : t \in [0, \omega]\}. \quad (1.5)$$

Moreover, (1.4) is fulfilled as an equality if and only if v is a constant function.

Proof. If v is a constant function, then (1.4) follows trivially.

Let, therefore, v be a non-constant function and choose $t_0, t_1 \in [0, \omega]$ such that

$$v(t_0) = M_v, \quad v(t_1) = m_v.$$

Without loss of generality we can assume that $t_0 < t_1$. Indeed, in the case where $t_1 < t_0$ we can consider a function $-v$ instead of v and using the fact that $v \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ we have

$$\int_0^\omega [v''(s)]_+ ds = \int_0^\omega [v''(s)]_- ds = \int_0^\omega [-v''(s)]_+ ds.$$

Put

$$M_1 = \max \{v'(t) : t \in [0, \omega]\}, \quad m_1 = \min \{v'(t) : t \in [0, \omega]\}.$$

Then, obviously, $M > 0$, $m < 0$ and by the periodicity of v and continuity of v' we have

$$M_v - m_v = \int_0^{t_0} v'(s) ds + \int_{t_1}^\omega v'(s) ds < M_1(t_0 + \omega - t_1) \quad (1.6)$$

and

$$M_v - m_v = - \int_{t_0}^{t_1} v'(s) ds < -m_1(t_1 - t_0). \quad (1.7)$$

On the other hand, we have $M_v - m_v \geq 0$ and thus the multiplying of the corresponding sides of (1.6) and (1.7) results in

$$(M_v - m_v)^2 < -m_1 M_1 (t_0 + \omega - t_1)(t_1 - t_0). \quad (1.8)$$

Now using the inequality $AB \leq \frac{1}{4}(A+B)^2$, from (1.8) we get

$$(M_v - m_v)^2 < \frac{(M_1 - m_1)^2 \omega^2}{16},$$

whence the inequality

$$M_v - m_v < \frac{\omega}{4}(M_1 - m_1) \quad (1.9)$$

follows.

On the other hand, choose $t_2, t_3 \in [0, \omega]$ such that

$$v'(t_2) = M_1, \quad v'(t_3) = m_1.$$

If $t_2 < t_3$ then by using again that v is ω -periodic we have

$$M_1 - m_1 = M_1 - v'(0) + v'(\omega) - m_1 = \int_0^{t_2} v''(s) ds + \int_{t_3}^\omega v''(s) ds \leq \int_0^\omega [v''(s)]_+ ds.$$

If $t_3 < t_2$ then

$$M_1 - m_1 = \int_{t_3}^{t_2} v''(s) ds \leq \int_0^\omega [v''(s)]_+ ds.$$

Consequently, in both cases $t_2 \leq t_3$ and $t_3 < t_2$ we have

$$M_1 - m_1 \leq \int_0^\omega [v''(s)]_+ ds$$

which together with (1.9) implies (1.4). □

2 The general case.

The following theorems are the main results of the paper.

Theorem 2.1. *Let $h \not\equiv 0$, $F > 0$, functions $w, \sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that the equalities*

$$w''(t) = Hg(t) - Gh(t) \quad \text{for a. e. } t \in [0, \omega], \quad (2.1)$$

$$\sigma''(t) = -\frac{F}{H}h(t) + f(t) \quad \text{for a. e. } t \in [0, \omega] \quad (2.2)$$

are fulfilled³ and let there exist $x_0 \in]0, +\infty[$ such that

$$x_0(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{H}{x_0GH + F}\right)^{1/\lambda} - \left(\frac{1}{x_0H}\right)^{1/\mu} \quad \text{for } t \in [0, \omega], \quad (2.3)$$

where

$$m_w = \min \{w(t) : t \in [0, \omega]\}, \quad m_\sigma = \min \{\sigma(t) : t \in [0, \omega]\}. \quad (2.4)$$

Then the problem (0.1), (0.2) has at least one positive solution.

Proof. Put

$$\alpha(t) = \left(\frac{1}{x_0H}\right)^{1/\mu} + x_0(w(t) - m_w) + \sigma(t) - m_\sigma \quad \text{for } t \in [0, \omega].$$

Then, obviously, $\alpha \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ and in view of (2.1) and (2.2) we have

$$\alpha''(t) = x_0Hg(t) - \left(x_0G + \frac{F}{H}\right)h(t) + f(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.5)$$

Moreover, according to (2.3) and (2.4),

$$\left(\frac{1}{x_0H}\right)^{1/\mu} \leq \alpha(t) \leq \left(\frac{H}{x_0GH + F}\right)^{1/\lambda} \quad \text{for } t \in [0, \omega]. \quad (2.6)$$

³see Remark 2.1 below.

Now (2.5) and (2.6) imply

$$\alpha''(t) \geq \frac{g(t)}{\alpha^\mu(t)} - \frac{h(t)}{\alpha^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega].$$

Consequently, α is a lower function to (0.1), (0.2).

Further, we can choose $x_1 \in]0, x_0]$ such that

$$x_1(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{1}{x_1 H}\right)^{1/\mu} - \left(\frac{H}{x_1 G H + F}\right)^{1/\lambda} \quad \text{for } t \in [0, \omega] \quad (2.7)$$

and put

$$\beta(t) = \left(\frac{H}{x_1 G H + F}\right)^{1/\lambda} + x_1(w(t) - m_w) + \sigma(t) - m_\sigma \quad \text{for } t \in [0, \omega].$$

Then, $\beta \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ and in view of (2.1) and (2.2) we have

$$\beta''(t) = x_1 H g(t) - \left(x_1 G + \frac{F}{H}\right) h(t) + f(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.8)$$

Moreover, according to (2.4) and (2.7),

$$\left(\frac{H}{x_1 G H + F}\right)^{1/\lambda} \leq \beta(t) \leq \left(\frac{1}{x_1 H}\right)^{1/\mu} \quad \text{for } t \in [0, \omega]. \quad (2.9)$$

Now (2.8) and (2.9) imply

$$\beta''(t) \leq \frac{g(t)}{\beta^\mu(t)} - \frac{h(t)}{\beta^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega].$$

Consequently, β is an upper function to (0.1), (0.2).

Moreover, (2.6) and (2.9) imply

$$\alpha(t) \leq \beta(t) \quad \text{for } t \in [0, \omega]. \quad (2.10)$$

Thus the assertion follows from Lemma 1.1. \square

Remark 2.1. Note that for every $q \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ such that $\int_0^\omega q(t)dt = 0$, the periodic solution v of the equation

$$v''(t) = q(t) \quad \text{for a. e. } t \in [0, \omega]$$

is given by the Green formula

$$v(t) = -\frac{1}{\omega} \left((\omega - t) \int_0^t s q(s) ds + t \int_t^\omega (\omega - s) q(s) ds \right) + c \quad \text{for } t \in [0, \omega], \quad (2.11)$$

where $c \in \mathbb{R}$. Therefore, the periodic functions w and σ with properties (2.1) and (2.2) exist and, moreover, are unique up to a constant term, the value of which has no influence on the validity of the condition (2.3). A similar observation can be made in relation to the formulations of the theorems given below.

Theorem 2.2. Let $\lambda > \mu$, $h \neq 0$, $g \neq 0$, $F = 0$, functions $w, \sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that the equalities (2.1) and

$$\sigma''(t) = f(t) \quad \text{for a. e. } t \in [0, \omega] \quad (2.12)$$

are fulfilled, and let there exist $x_0 \in]0, +\infty[$ such that

$$x_0(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{1}{x_0 G}\right)^{1/\lambda} - \left(\frac{1}{x_0 H}\right)^{1/\mu} \quad \text{for } t \in [0, \omega], \quad (2.13)$$

where m_w and m_σ are defined by (2.4). Then the problem (0.1), (0.2) has at least one positive solution.

Proof. Note that the inequality $\lambda > \mu$ implies

$$\lim_{x \rightarrow 0^+} \left(\frac{1}{xH}\right)^{1/\mu} - \left(\frac{1}{xG}\right)^{1/\lambda} = +\infty.$$

Therefore, analogously to the proof of Theorem 2.1, one can show that there exist lower and upper functions α, β satisfying (2.10). Consequently, the assertion follows from Lemma 1.1. \square

Corollary 2.1. Let $\lambda > \mu$, $h \neq 0$, $g \neq 0$, and let $w \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that (2.1) is fulfilled. Let, moreover,

$$M_w - m_w \leq \frac{H^{\frac{1+\lambda}{\lambda-\mu}}}{G^{\frac{1+\mu}{\lambda-\mu}}} \left(\frac{(1+\lambda)\mu}{(1+\mu)\lambda}\right)^{\frac{(1+\lambda)\mu}{\lambda-\mu}} \frac{\lambda - \mu}{(1+\mu)\lambda}, \quad (2.14)$$

where m_w is given by (2.4) and

$$M_w = \max \{w(t) : t \in [0, \omega]\}. \quad (2.15)$$

Then the problem (0.3), (0.2) has at least one positive solution.

Proof. In order to apply Theorem 2.2, put $f \equiv 0$, then $\sigma \equiv 0$. Take

$$x_0 = \left(\frac{(1+\mu)\lambda}{(1+\lambda)\mu}\right)^{\frac{\lambda\mu}{\lambda-\mu}} \frac{G^{\frac{\mu}{\lambda-\mu}}}{H^{\frac{\lambda}{\lambda-\mu}}}.$$

Then (2.14) implies (2.13), and thus the assertion follows from Theorem 2.2. \square

At this stage, Lemma 1.4 enables us to give a first concrete existence criterion.

Corollary 2.2. Let $\lambda > \mu$, $h \neq 0$, and $g \neq 0$. Let, moreover,

$$\frac{G^{1+\lambda}}{H^{1+\mu}} \leq \left(\frac{4}{\omega}\right)^{\lambda-\mu} \left(\frac{(1+\lambda)\mu}{(1+\mu)\lambda}\right)^{(1+\lambda)\mu} \left(\frac{\lambda - \mu}{(1+\mu)\lambda}\right)^{\lambda-\mu}. \quad (2.16)$$

Then the problem (0.3), (0.2) has at least one positive solution.

Proof. By Lemma 1.4, it is easy to verify that

$$M_w - m_w \leq \frac{\omega}{4}GH.$$

Now the assertion follows directly from Corollary 2.1. \square

To illustrate this latter result, we have selected a concrete physical model studied in [8, Section 5]. The dynamics of a trapless 3D Bose-Einstein condensate with variable scattering length is ruled by the equation

$$u''(t) = \frac{Q_1}{u^3} + \frac{a(t)Q_2}{u^4}, \quad (2.17)$$

where Q_1, Q_2 are positive parameters and $a(t)$ models the s-wave scattering length, which is assumed to vary ω -periodically in time. A negative $a(t)$ corresponds to attractive interactions between the elementary particles. Then the existence of a positive periodic solution of (2.17) is interpreted as a bound state of the condensate without external trap. Equation (2.17) is a particular case of (0.3) with $\mu = 3, \lambda = 4$. Then, a direct consequence of Corollary 2.2 is the existence of ω -periodic solution of (2.17) for any $a \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, $a(t) \leq 0$ for a. e. t , such that

$$\left(\int_0^\omega a(t)dt \right)^4 \geq \left(\frac{16}{15} \right)^{15} \frac{4Q_1^5\omega^6}{Q_2^4} \simeq 10.5315 \frac{Q_1^5\omega^6}{Q_2^4}.$$

The following results are devoted to the remaining cases $F < 0$ and $\mu > \lambda$. We are compelled to construct upper and lower functions on the reversed order.

Theorem 2.3. *Let $g \not\equiv 0, F < 0$, functions $w, \sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that the equalities (2.1) and*

$$\sigma''(t) = \frac{|F|}{G}g(t) + f(t) \quad \text{for a. e. } t \in [0, \omega] \quad (2.18)$$

are fulfilled, and let there exist $x_0 \in]0, +\infty[$ such that

$$x_0(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{G}{x_0GH + |F|} \right)^{1/\mu} - \left(\frac{1}{x_0G} \right)^{1/\lambda} \quad \text{for } t \in [0, \omega], \quad (2.19)$$

where m_w and m_σ are defined by (2.4). Moreover, let us define

$$\beta(t) = \left(\frac{1}{x_0G} \right)^{1/\lambda} + x_0(w(t) - m_w) + \sigma(t) - m_\sigma \quad \text{for } t \in [0, \omega] \quad (2.20)$$

and assume that $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$ verifies the property (P). Then the problem (0.1), (0.2) has at least one positive solution.

Proof. Put

$$\beta(t) = \left(\frac{1}{x_0 G} \right)^{1/\lambda} + x_0(w(t) - m_w) + \sigma(t) - m_\sigma \quad \text{for } t \in [0, \omega].$$

Then, $\beta \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ and in view of (2.1) and (2.18) we have

$$\beta''(t) = \left(x_0 H + \frac{|F|}{G} \right) g(t) - x_0 G h(t) + f(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.21)$$

Moreover, according to (2.4) and (2.19),

$$\left(\frac{1}{x_0 G} \right)^{1/\lambda} \leq \beta(t) \leq \left(\frac{G}{x_0 G H + |F|} \right)^{1/\mu} \quad \text{for } t \in [0, \omega]. \quad (2.22)$$

Now (2.21) and (2.22) imply

$$\beta''(t) \leq \frac{g(t)}{\beta^\mu(t)} - \frac{h(t)}{\beta^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega].$$

Consequently, β is an upper function to (0.1), (0.2).

Further, we can choose $x_1 \in]0, x_0]$ such that

$$x_1(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{1}{x_1 G} \right)^{1/\lambda} - \left(\frac{G}{x_1 G H + |F|} \right)^{1/\mu} \quad \text{for } t \in [0, \omega] \quad (2.23)$$

and put

$$\alpha(t) = \left(\frac{G}{x_1 G H + |F|} \right)^{1/\mu} + x_1(w(t) - m_w) + \sigma(t) - m_\sigma \quad \text{for } t \in [0, \omega].$$

Then, $\alpha \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ and in view of (2.1) and (2.18) we have

$$\alpha''(t) = \left(x_1 H + \frac{|F|}{G} \right) g(t) - x_1 G h(t) + f(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.24)$$

Moreover, according to (2.4) and (2.23),

$$\left(\frac{G}{x_1 G H + |F|} \right)^{1/\mu} \leq \alpha(t) \leq \left(\frac{1}{x_1 G} \right)^{1/\lambda} \quad \text{for } t \in [0, \omega]. \quad (2.25)$$

Now (2.24) and (2.25) imply

$$\alpha''(t) \geq \frac{g(t)}{\alpha^\mu(t)} - \frac{h(t)}{\alpha^\lambda(t)} + f(t) \quad \text{for a. e. } t \in [0, \omega].$$

Consequently, α is a lower function to (0.1), (0.2) and according to (2.22) and (2.25) we have

$$\beta(t) \leq \alpha(t) \quad \text{for } t \in [0, \omega]. \quad (2.26)$$

Furthermore, note that a function

$$\psi(y) = \frac{\mu}{\beta^{1+\mu}}y + \frac{1}{y^\mu}$$

is nondecreasing for $y \geq \beta$. Therefore we have

$$g(t) \left(\frac{\mu}{\beta^{1+\mu}(t)}u(t) + \frac{1}{u^\mu(t)} \right) - \frac{h(t)}{u^\lambda(t)} \leq g(t) \left(\frac{\mu}{\beta^{1+\mu}(t)}v(t) + \frac{1}{v^\mu(t)} \right) - \frac{h(t)}{v^\lambda(t)} \quad \text{for } t \in [0, \omega]$$

whenever $\beta(t) \leq u(t) \leq v(t)$ for $t \in [0, \omega]$, whence we get

$$\frac{g(t)}{u^\mu(t)} - \frac{h(t)}{u^\lambda(t)} - \left(\frac{g(t)}{v^\mu(t)} - \frac{h(t)}{v^\lambda(t)} \right) \leq \frac{\mu g(t)}{\beta^{1+\mu}(t)}(v(t) - u(t)).$$

Thus the assertion follows from Lemma 1.2. \square

Theorem 2.4. *Let $\mu > \lambda$, $h \not\equiv 0$, $g \not\equiv 0$, $F = 0$, functions $w, \sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that the equalities (2.1) and (2.12) are fulfilled, and let there exist $x_0 \in]0, +\infty[$ such that*

$$x_0(w(t) - m_w) + \sigma(t) - m_\sigma \leq \left(\frac{1}{x_0 H} \right)^{1/\mu} - \left(\frac{1}{x_0 G} \right)^{1/\lambda} \quad \text{for } t \in [0, \omega], \quad (2.27)$$

where m_w and m_σ are defined by (2.4). Moreover, assume that $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$ verifies the property (P), where β is given by (2.20). Then the problem (0.1), (0.2) has at least one positive solution.

Proof. Note that the inequality $\mu > \lambda$ implies

$$\lim_{x \rightarrow 0^+} \left(\frac{1}{xG} \right)^{1/\lambda} - \left(\frac{1}{xH} \right)^{1/\mu} = +\infty.$$

Therefore, analogously to the proof of Theorem 2.3, one can show that there exist lower and upper functions α, β satisfying (2.26). Consequently, the assertion follows from Lemma 1.2 with $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$. \square

Corollary 2.3. *Let $\mu > \lambda$, $h \not\equiv 0$, $g \not\equiv 0$, $w \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that (2.1) is fulfilled, and let*

$$M_w - m_w \leq \frac{G^{\frac{1+\mu}{\mu-\lambda}}}{H^{\frac{1+\lambda}{\mu-\lambda}}} \left(\frac{(1+\mu)\lambda}{(1+\lambda)\mu} \right)^{\frac{(1+\mu)\lambda}{\mu-\lambda}} \frac{\mu - \lambda}{(1+\lambda)\mu}, \quad (2.28)$$

where m_w and M_w are given by (2.4) and (2.15), respectively. Moreover, let us define

$$\beta(t) = \left(\frac{(1+\mu)\lambda}{(1+\lambda)\mu} \right)^{\frac{\mu}{\mu-\lambda}} \left(\frac{G}{H} \right)^{\frac{1}{\mu-\lambda}} + \left(\frac{(1+\lambda)\mu}{(1+\mu)\lambda} \right)^{\frac{\lambda\mu}{\mu-\lambda}} \frac{H^{\frac{\lambda}{\mu-\lambda}}}{G^{\frac{\mu}{\mu-\lambda}}} (w(t) - m_w) \quad \text{for } t \in [0, \omega] \quad (2.29)$$

and assume that $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$ verifies the property (P). Then the problem (0.3), (0.2) has at least one positive solution.

Proof. Put $f \equiv 0$ and

$$x_0 = \left(\frac{(1+\lambda)\mu}{(1+\mu)\lambda} \right)^{\frac{\lambda\mu}{\mu-\lambda}} \frac{H^{\frac{\lambda}{\mu-\lambda}}}{G^{\frac{\mu}{\mu-\lambda}}}.$$

Then the assertion follows from Theorem 2.4. \square

Corollary 2.4. Let $\mu > \lambda$, $h \neq 0$, and $g \neq 0$. Let, moreover,

$$\frac{H^{1+\mu}}{G^{1+\lambda}} \leq \left(\frac{4}{\omega} \right)^{\mu-\lambda} \left(\frac{(1+\mu)\lambda}{(1+\lambda)\mu} \right)^{(1+\mu)\lambda} \left(\frac{\mu-\lambda}{(1+\lambda)\mu} \right)^{\mu-\lambda} \times \min \left\{ 1, \left(\frac{1+\lambda}{\mu-\lambda} \right)^{\mu-\lambda} \left(\frac{(1+\mu)\lambda}{(1+\lambda)\mu} \right)^{(\mu-\lambda)(1+\mu)} \right\}. \quad (2.30)$$

Then the problem (0.3), (0.2) has at least one solution.

Proof. According to Lemma 1.4, the inequality (2.30) implies (2.28) and moreover, after some tedious computations one has

$$\int_0^\omega \varphi(s) ds = \mu \int_0^\omega \frac{g(s)}{\beta^{1+\mu}(s)} ds \leq \frac{4}{\omega},$$

with β defined by (2.29). Consequently, by Lemma 1.3, $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$ verifies the property (P) and the assertion follows from Corollary 2.3. \square

To finish this section, we remark that our approach does not cover the case $\lambda = \mu$, $F = 0$ which is of particular interest for applications (see the introduction of [1]). The following problem is unsolved.

Open problem 2.1. If $\lambda = \mu$, we know that $H > G > 0$ is a necessary condition for the existence of a positive solution of problem (0.3) (0.2). Prove that it is also sufficient.

3 The attractive case.

In this section we focus on the equation with a pure attractive singularity, that is, the case when $g \equiv 0$.

Corollary 3.1. *Let $h \not\equiv 0$, $F > 0$, $\sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that (2.2) is fulfilled, and let*

$$(M_\sigma - m_\sigma)^\lambda F < H, \quad (3.1)$$

where m_σ is defined by (2.4) and

$$M_\sigma = \max \{ \sigma(t) : t \in [0, \omega] \}. \quad (3.2)$$

Then the problem (0.4), (0.2) has at least one positive solution.

Proof. The assertion follows from Theorem 2.1 with $g \equiv 0$. □

Corollary 3.2. *Let $h \not\equiv 0$, $F > 0$, and let*

$$\left(\frac{\omega}{4} F_+ \right)^\lambda F \leq H. \quad (3.3)$$

Then the problem (0.4), (0.2) has at least one positive solution.

Proof. By Lemma 1.4, in view of $F > 0$, we have

$$M_\sigma - m_\sigma < \frac{\omega}{4} F_+.$$

Now the assertion follows from Corollary 3.1 in a trivial way. □

The latter result is new even for the original equation posed by Lazer and Solimini,

$$u''(t) = -\frac{1}{u^\lambda(t)} + f(t). \quad (3.4)$$

In [7], it is proved that if f is continuous and ω -periodic, then $F > 0$ is a necessary and sufficient condition for the existence of a positive ω -periodic solution. Here we are extending partially this result to the case when $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$. On the other hand, even if f is continuous, then $F > 0$ is not sufficient condition for the existence of a positive ω -periodic solution to the equation (0.4) in the case, when h is possibly zero on the set of a positive measure, as shown in the following example.

Counter-example 3.1. Let $\varepsilon \in]0, \omega/4[$ and put

$$f(t) = \begin{cases} 0 & \text{for } t \in [0, \frac{\omega}{2} - \varepsilon[\cup]\frac{\omega}{2} + \varepsilon, \omega] \\ \frac{2}{\varepsilon}(t - \frac{\omega}{2} + \varepsilon) & \text{for } t \in]\frac{\omega}{2} - \varepsilon, \frac{\omega}{2}[\\ \frac{2}{\varepsilon}(\frac{\omega}{2} + \varepsilon - t) & \text{for } t \in [\frac{\omega}{2}, \frac{\omega}{2} + \varepsilon[\end{cases},$$

$$h(t) = \begin{cases} -\frac{t^2}{2} + \varepsilon(\frac{\omega}{2} - \varepsilon) & \text{for } t \in [0, \varepsilon[\\ 0 & \text{for } t \in [\varepsilon, \omega - \varepsilon[\\ -\frac{(\omega-t)^2}{2} + \varepsilon(\frac{\omega}{2} - \varepsilon) & \text{for } t \in [\omega - \varepsilon, \omega] \end{cases},$$

$$v(t) = \begin{cases} -\frac{t^2}{2} + \varepsilon(\frac{\omega}{2} - \varepsilon) & \text{for } t \in [0, \varepsilon[\\ \varepsilon(\frac{\omega}{2} - t) - \frac{\varepsilon^2}{2} & \text{for } t \in [\varepsilon, \frac{\omega}{2} - \varepsilon[\\ \frac{(t - \frac{\omega}{2} + \varepsilon)^3}{3\varepsilon} + \varepsilon(\frac{\omega}{2} - t) - \frac{\varepsilon^2}{2} & \text{for } t \in [\frac{\omega}{2} - \varepsilon, \frac{\omega}{2}[\\ \frac{(\frac{\omega}{2} + \varepsilon - t)^3}{3\varepsilon} + \varepsilon(t - \frac{\omega}{2}) - \frac{\varepsilon^2}{2} & \text{for } t \in [\frac{\omega}{2}, \frac{\omega}{2} + \varepsilon[\\ \varepsilon(t - \frac{\omega}{2}) - \frac{\varepsilon^2}{2} & \text{for } t \in [\frac{\omega}{2} + \varepsilon, \omega - \varepsilon[\\ -\frac{(\omega-t)^2}{2} + \varepsilon(\frac{\omega}{2} - \varepsilon) & \text{for } t \in [\omega - \varepsilon, \omega] \end{cases},$$

and

$$\sigma(t) = -\frac{1}{\omega} \left[(\omega - t) \int_0^t s \left(-\frac{F}{H} h(s) + f(s) \right) ds + t \int_t^\omega (\omega - s) \left(-\frac{F}{H} h(s) + f(s) \right) ds \right],$$

where

$$H = \int_0^\omega h(s) ds = 2\varepsilon^2 \left(\frac{\omega}{2} - \varepsilon \right) - \frac{\varepsilon^3}{3}, \quad F = F_+ = \int_0^\omega f(s) ds = 2\varepsilon.$$

Obviously, f is continuous, $v, \sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, $\sigma(t) = \sigma(\omega - t)$ for $t \in [0, \omega]$, and consequently, $\sigma'(t) = -\sigma'(\omega - t)$ for $t \in [0, \omega]$. Therefore,

$$\sigma'(\omega) = \sigma'(0) = -\sigma'(\omega), \quad \sigma'(\omega/2) = -\sigma'(\omega/2),$$

which implies $\sigma'(0) = 0$, $\sigma'(\omega/2) = 0$. Moreover, now it can be easily verified that

$$\max \{ \sigma(t) : t \in [0, \omega] \} = \sigma(0), \quad \min \{ \sigma(t) : t \in [0, \omega] \} = \sigma(\omega/2).$$

Thus

$$M_\sigma - m_\sigma = \sigma(0) - \sigma(\omega/2) = \varepsilon \left(\frac{\omega}{2} - \varepsilon \right) + \frac{\varepsilon^2}{6} + \frac{\varepsilon^3}{12\omega - 28\varepsilon},$$

$$\frac{H}{F} = \varepsilon \left(\frac{\omega}{2} - \varepsilon \right) - \frac{\varepsilon^2}{6}, \quad \frac{\omega}{4} F_+ = \frac{\omega}{2} \varepsilon.$$

We will show that the problem

$$u'' = -\frac{h(t)}{u} + f(t); \quad u(0) = u(\omega), \quad u'(0) = u'(\omega) \quad (3.5)$$

has no positive solution. Suppose on the contrary, that there exists a positive solution u to (3.5). Put $w(t) = v(t) - u(t)$ for $t \in [0, \omega]$. Then

$$w'' = p(t)w; \quad w(0) = w(\omega), \quad w'(0) = w'(\omega)$$

with

$$p(t) = \begin{cases} \frac{1}{u(t)} & \text{for } t \in [0, \varepsilon[\cup [\omega - \varepsilon, \omega] \\ 0 & \text{for } t \in [\varepsilon, \omega - \varepsilon[\end{cases}.$$

Consequently $w \equiv 0$, i.e. $u \equiv v$. However, $v(\omega/2) = -\varepsilon^2/6 < 0$, which contradicts our assumption.

This example shows that the inequalities (3.1) and (3.3) in Corollaries 3.1 and 3.2 are optimal in a certain sense and cannot be improved. In particular, the condition (3.1), resp. (3.3), cannot be replaced by the condition

$$(M_\sigma - m_\sigma)^\lambda F \leq H + \varepsilon,$$

resp.

$$\left(\frac{\omega}{4}F_+\right)^\lambda F \leq H + \varepsilon,$$

no matter how small ε is.

We finish the section with two open questions.

Open problem 3.1. Let us assume $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, and $\lambda > 0$. Prove or disprove that $F > 0$ is a necessary and sufficient condition for the existence of a ω -periodic positive solution of the Lazer-Solimini equation (3.4).

Open problem 3.2. Let us assume $h \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$, $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, $\lambda > 0$, and

$$\text{meas} \{t \in [0, \omega] : h(t) = 0\} = 0.$$

Find a condition different from (3.1) (resp. (3.3)) sufficient for the existence of a positive solution of problem (0.4), (0.2).

4 The repulsive case.

Finally, we analyze the equation with a pure repulsive singularity, that is, the case when $h \equiv 0$.

Corollary 4.1. Let $g \not\equiv 0$, $F < 0$, $\sigma \in AC^1(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ be such that (2.18) is fulfilled, and let

$$(M_\sigma - m_\sigma)^\mu |F| < G, \tag{4.1}$$

where m_σ and M_σ is defined by (2.4) and (3.2), respectively. Let, moreover, either

$$\frac{\mu |F|^{\frac{1+\mu}{\mu}} g(t)}{(G^{1/\mu} - |F|^{1/\mu} (M_\sigma - \sigma(t)))^{1+\mu}} \leq \left(\frac{\pi}{\omega}\right)^2 \quad \text{for a. e. } t \in [0, \omega], \tag{4.2}$$

or

$$\mu|F|^{\frac{1+\mu}{\mu}} \int_0^\omega \frac{g(s)}{(G^{1/\mu} - |F|^{1/\mu}(M_\sigma - \sigma(s)))^{1+\mu}} ds \leq \frac{4}{\omega}. \quad (4.3)$$

Then the problem (0.5), (0.2) has at least one positive solution.

Proof. Put $h \equiv 0$,

$$x_0 = \frac{|F|^{\lambda/\mu}}{G(G^{1/\mu} - |F|^{1/\mu}(M_\sigma - m_\sigma))^\lambda},$$

and define a function β by (2.20). After some algebra,

$$\beta(t) = \left(\frac{G}{|F|} \right)^{1/\mu} - M_\sigma + \sigma(t) \quad \text{for } t \in [0, \omega]$$

and each of (4.2) and (4.3) guarantees that $\varphi(t) = \frac{\mu g(t)}{\beta^{1+\mu}(t)}$ satisfies the property (P). Moreover, (4.1) yields (2.19). Therefore the assertion follows from Theorem 2.3. \square

Corollary 4.2. *Let $g \not\equiv 0$ and $F < 0$. Let, moreover,*

$$\left(\frac{\omega}{4} \mu G \right)^{\frac{1}{1+\mu}} |F|^{1/\mu} + \frac{\omega}{4} F_- |F|^{1/\mu} \leq G^{1/\mu}. \quad (4.4)$$

Then the problem (0.5), (0.2) has at least one positive solution.

Proof. According to Lemma 1.4,

$$M_\sigma - m_\sigma \leq \frac{\omega}{4} F_-.$$

Then, the inequality (4.4) implies both (4.1) and (4.3). Consequently, the assertion follows from Corollary 4.1. \square

Again, this result is new even for the original equation posed by Lazer and Solimini,

$$u''(t) = \frac{1}{u^\mu(t)} + f(t). \quad (4.5)$$

In [7], it is proved that if $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ and $\mu \geq 1$ (strong force assumption), then $F < 0$ is a necessary and sufficient condition for the existence of a positive ω -periodic solution. Moreover, it is shown with a counterexample that the strong force assumption cannot be dropped without additional conditions. Later, in [11] the authors proved that (4.5) with $\mu < 1$ has a positive ω -periodic solution if $F < 0$ and

$$f(t) \geq - \left(\frac{\pi^2}{\omega^2 \mu} \right)^{\frac{\mu}{\mu+1}} (\mu + 1) \quad \text{for a. e. } t \in [0, \omega].$$

Therefore, a uniform bound from below is required. The importance of Corollary 4.2 in this context relies in that it provides for the first time a sufficient existence condition for a truly $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$ (possibly unbounded). Of course, the main question remains open.

Open problem 4.1. Let us assume $g \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R}_+)$, $g \not\equiv 0$, $f \in L(\mathbb{R}/\omega\mathbb{Z}; \mathbb{R})$, and $\mu > 0$. Find a necessary and sufficient condition over f, g for the existence of a positive solution of problem (0.5), (0.2).

References

- [1] J. L. Bravo, P. J. Torres, *Periodic solutions of a singular equation with indefinite weight*, preprint.
- [2] C. De Coster and P. Habets, *Upper and lower solutions in the theory of ODE boundary value problems: classical and recent results*, in: *Nonlinear Analysis and Boundary Value Problems for Ordinary Differential Equations*, edited by F. Zanolin, CISM–ICMS 371 (Springer–Verlag, New York, 1996), 1–78.
- [3] P. Habets, L. Sanchez, *Periodic solutions of some Linard equations with singularities*, Proc. Amer. Math. Soc. 109 (1990), No. 4, 1035–1044.
- [4] K. Johansson, *On Separation of Phases in One-Dimensional Gases*, Commun. Math. Phys. **169**, (1995), 521-561.
- [5] J. Lei, M. Zhang, *Twist property of periodic motion of an atom near a charged wire*, Lett. Math. Phys. 60 (2002), No. 1, 9–17.
- [6] M. Kunze, *Periodic solutions of a singular Lagrangian system related to dispersion-managed fiber communication devices*, Nonlinear Dynamics and Systems Theory 1, (2001), 159-167.
- [7] A. C. Lazer, S. Solimini, *On periodic solutions of nonlinear differential equations with singularities*, Proc. Am. Math. Soc. **99** (1987), No. 1, 109–114.
- [8] G. D. Montesinos, V. M. Perez-García, P. J. Torres, *Stabilization of solitons of the multidimensional nonlinear Schrodinger equation: matter-wave breathers*, Physica D **191** (2004), 193-210.
- [9] H. N. Pishkenari, M. Behzad, A. Meghdari, *Nonlinear dynamic analysis of atomic force microscopy under deterministic and random excitation*, Chaos, Solitons & Fractals, **37**, Iss. 3 (2008), 748-762.
- [10] I. Rachůnková, S. Staněk, M. Tvrdý, *Singularities and Laplacians in Boundary Value Problems for Nonlinear Ordinary Differential Equations*, in Handbook of Differential Equations (Ordinary Differential Equations) vol. 3, Elsevier (2006).
- [11] I. Rachůnková, M. Tvrdý, I. Vrkoč, *Existence of nonnegative and nonpositive solutions for second order periodic boundary value problems*, J. Differential Equations 176 (2001), 445-469.
- [12] S. Rützel, S. I. Lee, A. Raman, *Nonlinear dynamics of atomic-force-microscope probes driven in Lennard-Jones potentials*, Proc. R. Soc. Lond. A 2003 459, 1925-1948.
- [13] P. J. Torres, *Existence and stability of periodic solutions for second order semilinear differential equations with a singular nonlinearity*, Proceedings of the Royal Society of Edinburgh **137A** (2007), 195-201.

- [14] P. J. Torres, M. Zhang, *A monotone iterative scheme for a nonlinear second order equation based on a generalized anti-maximum principle*, Math. Nachr. **251** (2003) / DOI 10.1002/mana.200310033, 101-107.
- [15] G. Yang, J. Lu, A.C.J. Luo, *On the computation of Lyapunov exponents for forced vibration of a Lennard-Jones oscillator*, Chaos, Solitons & Fractals **23** (2005), 833-841.