

Differential
Equations
& Applications

*With Compliments
of the Author*

Zagreb, Croatia

Volume 3, Number 3, August 2011

Alexander Gutiérrez and Pedro J. Torres

*Periodic solutions of Liénard equation with one or
two weak singularities*

DIFFERENTIAL EQUATIONS & APPLICATIONS

AIMS AND SCOPE

Differential Equations and Applications (DEA, Differ. Equ. Appl.) will publish original papers from all branches of ordinary, functional-differential, and partial differential equations. Only papers of the highest quality will be accepted for the publication. The papers which demonstrate novelty or give relations of differential equations with other fields of mathematics and applications are preferred.

DEA is published quarterly, in February, May, August and November.

SUBMISSION

Manuscripts should be submitted electronically in PostScript or Adobe Acrobat PDF format to the DEA Editorial Office through DEA page www.ele-math.com or by e-mail dea@ele-math.com

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the DEA regarding the article. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to DEA. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal, and that its submission for publication has been approved by all the authors.

PREPARATION OF MANUSCRIPT

Manuscripts should be written in English.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send final version of file(s) to the Editorial Office.

TITLE PAGE

A title page should include the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author and running head less than 60 characters. Authors should be listed alphabetically.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2000) with primary (and secondary) subject classification codes and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The author should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS

The author will receive proofs only once. Late return of the proofs will delay the article to a later issue.

PAGE CHARGES, OFFPRINTS

There are no page charges. Authors jointly receive 30 offprints of their articles free of charge. Additional offprints may be ordered from DEA prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

DEA is published by Publishing House **ELEMENT**, Zagreb, Croatia, and
Department of Applied Mathematics, FER, University of Zagreb.

All correspondence and subscription orders should be addressed to the Editorial Office:

Differential Equations & Applications
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia

www.ele-math.com
e-mail: dea@ele-math.com
Fax: +385 1 6008799

PERIODIC SOLUTIONS OF LIÉNARD EQUATION WITH ONE OR TWO WEAK SINGULARITIES

ALEXANDER GUTIÉRREZ AND PEDRO J. TORRES

(Communicated by V. Županović)

Abstract. In this paper we study the existence and asymptotic stability of periodic solutions of the differential equation

$$\ddot{x} + f(x)\dot{x} + g(x) = h(t),$$

where $h(t)$ is T -periodic, $f(x)$ is positive and $g(x)$ is strictly monotonically increasing and has one or two weak singularities. The method of proof relies on the construction of a positively invariant region of the flux.

1. Introduction

In this paper we deal with the existence and stability of T -periodic solutions of a second order differential equation of Liénard-type

$$\ddot{x} + f(x)\dot{x} + g(x) = h(t), \quad (1)$$

where $h(t)$ is a continuous and T -periodic function, $f, g: (l_1, l_2) \rightarrow \mathbb{R}$ are locally Lipschitz continuous functions.

In principle, $-\infty \leq l_1 < l_2 \leq +\infty$, but we are interested in the case where at least one of them is finite.

The study of scalar second order equations with singularities can be traced back to a paper by Nagumo [10] published in 1944. It is important to remark this fact because it seems to be little known, and up to our knowledge it is not recorded in the related literature. The available reviews [9, 12, 13] register as early references some papers by Forbat, Huaux and Derwindé in the sixties [4, 5, 2]. Although the first application of topological degree is due to Fauré [3], the papers [7, 6] constitute the landmarks on this topic.

In the study of equations with singularities, the so-called strong force assumption has played a prominent role. To explain it, let us define \bar{x} as the minimum of the values of (l_1, l_2) such that $g(\bar{x}) = 0$ and let us define the potential

$$G(x) := \int_{\bar{x}}^x g(s) ds.$$

Mathematics subject classification (2010): 34C25.

Keywords and phrases: periodic solution, singularity, asymptotic stability.

Supported by projects MTM2005-03483 and FQM2216.

Then, it is said that g has a strong singularity at l_1 (resp. l_2) if

$$\lim_{x \rightarrow l_1^+} G(x) = +\infty, \quad (\text{resp. } \lim_{x \rightarrow l_2^-} G(x) = +\infty).$$

On the other hand, when such a limit is finite, we speak about a weak singularity. All the classical papers mentioned before assume a strong force condition. In fact, in the seminal paper of Lazer and Solimini [7] it is shown that such a condition can not be dropped without further assumptions. On this basis, such a condition became standard in the related works. However, in the latter years the interest on weak singularities has increased and some conditions for existence of periodic solutions [1, 8, 11, 14, 15, 16, 17, 18] can be found. Our purpose in this paper is to recover the original method of Nagumo developed in [10] for strong singularities and show that it is suitable to deal also with weak singularities.

In order to formulate our main result, let us define the following functions

$$\begin{aligned} F(x) &:= \int_x^x f(s) ds, \\ W(x) &:= \frac{F^2(x)}{4} + 2G(x), \\ Z(x) &:= \frac{F^2(x)}{2} + 2G(x). \end{aligned}$$

In the following, $\|\cdot\|_\infty$ stands for the usual supremum norm.

THEOREM 1. *Let us assume the following hypotheses.*

- (H1) $C^* := \inf\{W(l_1), W(l_2)\} < \infty$.
- (H2) *There exists $f_0 > 0$ such that $f(x) \geq f_0$ for every $x \in (l_1, l_2)$.*
- (H3) *$g(x)$ is a strictly increasing function that changes sign.*

Fix $0 < C < C^$ and let $l'_1 < l'_2$ be the solutions of $Z(x) = C$. Define the function*

$$R(x) := -\frac{1}{2}|F(x)| + \sqrt{C^* - W(x)}, \tag{2}$$

and fix the following positive constants

$$\begin{aligned} K_1 &:= \min_{i=1,2} \left\{ \frac{|g(l'_i)|}{f_0}, \sqrt{\frac{2g(l'_i)F(l'_i)}{f_0}} \right\}, \\ K_2 &:= \min_{i=1,2} \left\{ \frac{1}{2}R(l'_i), \frac{1}{|F(l'_i)|}R^2(l'_i) \right\}. \end{aligned}$$

Then, under the assumption

$$\|h\|_\infty < \frac{f_0}{2} \min\{K_1, K_2\}, \tag{3}$$

there exists at least one T -periodic solution $\varphi(t)$ of Eq. (1). Such solution verifies

$$l'_1 \leq \varphi(t) \leq l'_2 \quad \text{for all } t. \tag{4}$$

Some comments are pertinent here. Condition (H1) is a weak force assumption. (H2) is a dissipative condition. Finally, (H3) implies that the statement is consistent. In fact, if $g(x)$ and $f(x)$ satisfy (H2)-(H3) then functions $W(x)$, $Z(x)$ satisfy

$$\begin{aligned} W'(x), Z'(x) < 0, \quad x \in (l_1, \bar{x}), \\ W'(x), Z'(x) > 0, \quad x \in (\bar{x}, l_2). \end{aligned}$$

Hence $Z(x)$ and $W(x)$ have an absolute minimum at \bar{x} and $W(\bar{x}) = Z(\bar{x}) = 0$. Moreover, for all $0 < C \leq C^*$ the equation $Z(x) = C$ has exactly two solutions l'_1, l'_2 with $l'_1 < \bar{x} < l'_2$. Finally, the function $R(x)$ is positive for $x \in (l'_1, l'_2)$ (see Lemma 1) and therefore the constant K_2 is in fact positive.

The paper is structured into four section. After this Introduction, Section 2 is devoted to prove the main result. In Section 3 we show how to take advantage on the bounds obtained in the previous section in order to get a result on asymptotic stability. Finally, in Section 4 we provide some illustrative examples and compare our results with those available in the related literature.

2. Proof of the main result

We begin with a preliminary lemma.

LEMMA 1. *The function $R(x)$ is positive for $x \in (l'_1, l'_2)$.*

Proof. If $x \in (l'_1, l'_2)$, we have

$$Z(x) = W(x) + \frac{1}{4}F^2(x) < C \text{ and } C^* - W(x) > C^* - C + \frac{1}{4}F^2(x) > 0.$$

Then

$$\sqrt{C^* - W(x)} > \sqrt{C^* - C + \frac{1}{4}F^2(x)} > \frac{1}{2}|F(x)|. \quad \square$$

With this lemma, $K_2 > 0$ and the assumptions of Theorem 1 are consistent. Now, let us rewrite (1) as a system

$$\begin{cases} \dot{x} = y - F(x), \\ \dot{y} = -g(x) + h(t). \end{cases} \tag{5}$$

The proof of Theorem 1 will consist of establishing a positively invariant region for system (5). To this purpose, let us define the energy functional

$$P(x,y) := \left(y - \frac{F(x)}{2} \right)^2 + W(x). \tag{6}$$

Obviously, the inequality $P(x,y) \leq C^*$ is equivalent to

$$\frac{F(x)}{2} - \sqrt{C^* - W(x)} \leq y \leq \frac{F(x)}{2} + \sqrt{C^* - W(x)}. \tag{7}$$

Moreover, it is easy to realize that $P(x, y) = C^*$ is a simple closed curve. The goal is to prove that the simply connected set defined by

$$D := \{(x, y) : l'_1 \leq x \leq l'_2, P(x, y) \leq C^*\}, \tag{8}$$

is positively invariant for system (5).

The following auxiliary result will be useful. For convenience, in the rest of the section we will call $H = \|h\|_\infty$.

LEMMA 2. *Under the conditions of Theorem 1, if $x \in (l'_1, l'_2)$ and $P(x, y) \geq C^*$ then the following inequalities are fulfilled*

$$|y - F(x)| \geq \frac{4H}{f_0}, \tag{9}$$

$$(y - F(x))^2 > \frac{2H}{f_0} |F(x)|. \tag{10}$$

Proof. In order to prove (9), we must meet that

$$y > \frac{4H}{f_0} + F(x) \quad \text{or} \quad y < -\frac{4H}{f_0} + F(x).$$

Since $P(x, y) \geq C^*$, from (7) we have

$$y \geq \frac{F(x)}{2} + \sqrt{C^* - W(x)},$$

or

$$y \leq \frac{F(x)}{2} - \sqrt{C^* - W(x)}.$$

Therefore, it is sufficient to prove

$$\begin{aligned} F(x) + \frac{4H}{f_0} &< \frac{F(x)}{2} + \sqrt{C^* - W(x)}, \\ F(x) - \frac{4H}{f_0} &> \frac{F(x)}{2} - \sqrt{C^* - W(x)}, \end{aligned}$$

for all $x \in (l'_1, l'_2)$, that is,

$$H < \frac{f_0}{4} \left(-\frac{|F(x)|}{2} + \sqrt{C^* - W(x)} \right) = \frac{f_0}{4} R(x). \tag{11}$$

Note that from the definition of l'_1, l'_2 we have

$$\sqrt{C^* - W(x)} \geq \frac{1}{2} |F(x)| \quad \text{for all } x \in (l'_1, l'_2).$$

On the other hand, it is easy to verify that $R'(x) > 0$ for $x \in (l'_1, \bar{x})$ and $R'(x) < 0$ for $x \in (\bar{x}, l'_2)$. Therefore

$$\min_{x \in [l'_1, l'_2]} R(x) = \min_{i=1,2} R(l'_i).$$

By using the condition (3),

$$H < \frac{f_0}{2}K_2 \leq \frac{f_0}{4} \min_{i=1,2} R(l'_i) \leq \frac{f_0}{4}R(x),$$

which is just inequality (11), thus (9) is proved.

Similarly we have that H satisfies (10) if

$$0 < H < \frac{f_0}{2|F(x)|} \left(-\frac{|F(x)|}{2} + \sqrt{C^* - W(x)} \right)^2 = \frac{f_0}{2|F(x)|}R^2(x). \tag{12}$$

Note that

$$\begin{aligned} \frac{d}{dx} \left(\frac{1}{|F(x)|}R^2(x) \right) &> 0, \quad x \in (l'_1, \bar{x}), \\ \frac{d}{dx} \left(\frac{1}{|F(x)|}R^2(x) \right) &< 0, \quad x \in (\bar{x}, l'_2). \end{aligned}$$

Therefore

$$\min_{x \in [l'_1, l'_2]} \frac{R^2(x)}{|F(x)|} = \min_{i=1,2} \frac{R^2(l'_i)}{|F(l'_i)|}.$$

By using (3),

$$H < \frac{f_0}{2}K_2 \leq \frac{f_0}{2} \frac{R^2(l'_i)}{|F(l'_i)|} \leq \frac{f_0}{2} \frac{R^2(x)}{|F(x)|},$$

which is just inequality (12). \square

Now, we are ready to prove the main theorem.

PROOF OF THEOREM 1. We are going to prove that the region D defined by (8) is positively invariant. It is sufficient to prove that

$$\dot{P}(x,y) < 0, \quad (x,y) \notin D, \tag{13}$$

where \dot{P} is the total derivative of P .

Taking into account (6) and (5),

$$\begin{aligned} \dot{P} &= [y + (y - F(x))] \dot{y} + [-f(x)(y - F(x)) + 2g(x)] \dot{x} \\ &= [2y - F(x)](h(t) - g(x)) + [2g(x) - f(x)(y - F(x))](y - F(x)) \\ &= h(t)[2y - F(x)] - g(x)[2y - F(x)] - f(x)(y - F(x))^2 + 2g(x)(y - F(x)) \\ &= -f(x)(y - F(x))^2 + 2h(t)(y - F(x)) - F(x)g(x) + F(x)h(t). \end{aligned} \tag{14}$$

We distinguish two cases (see Figure 1).

Case 1. Let be $x \in (l_1, l'_1] \cup [l'_2, l_2)$. Note $F(x)g(x)$ and $|g(x)|$ are nonnegative functions with an absolute minimum in \bar{x} and such that they are decreasing in $(l_1, l'_1]$




Figure 1: The white region D is positively invariant for system (5)

and increasing in $[l'_1, l_2)$. By (3), we have that $H < \frac{f_0}{2}K_1$, and considering the definition of K_1 and the previous argument, the following inequalities hold

$$F(x)g(x) > \frac{2H^2}{f_0}, \quad |g(x)| > 2H$$

for all $x \in (l_1, l'_1] \cup [l'_2, l_2)$. In consequence,

$$\begin{aligned} -f(x)(y - F(x))^2 &\leq -f_0(y - F(x))^2, \\ 2h(x)(y - F(x)) &\leq 2H|y - F(x)|, \end{aligned}$$

and

$$\begin{aligned} F(x)h(x) - F(x)g(x) &< |F(x)|H - F(x)g(x) \\ &< |F(x)|\frac{|g(x)|}{2} - F(x)g(x) \\ &\leq -\frac{1}{2}F(x)g(x) < -\frac{H^2}{f_0}. \end{aligned}$$

Putting the above inequalities into (14), it follows

$$\dot{P} < -\left(\sqrt{f_0}|y - F(x)| - \frac{H}{\sqrt{f_0}}\right)^2 \leq 0,$$

for all $x \in (l_1, l'_1] \cup [l'_2, l_2)$ and $y \in \mathbb{R}$.

Case 2. Let be $x \in (l'_1, l'_2)$ and $P(x, y) \geq C^*$. By Lemma 2, the inequalities (9)-(10) are satisfied and therefore

$$\begin{aligned}
 -f(x)(y - F(x))^2 &\leq -f_0(y - F(x))^2, \\
 F(x)h(t) - F(x)g(x) &\leq H|F(x)| < \frac{f_0}{2}(y - F(x))^2.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 \dot{P} &< |y - F(x)| \left(-\frac{f_0}{2}|y - F(x)| + 2h(t) \right) \\
 &< |y - F(x)| (-2H + 2h(t)) < 0,
 \end{aligned}$$

for all $x \in (l'_1, l'_2)$ and $y \in \mathbb{R}$ such that $P(x, y) \geq C^*$.

Therefore, combining Case 1 and Case 2 we get (13).

Finally using (13), equation (1) has a T -periodic solution by a basic application of Brouwer's fixed point theorem. Denote by $(x(t; t_0, x_0, y_0), y(t; t_0, x_0, y_0))$ the unique solution of the Cauchy problem for system (5). Using the fact that D is positively invariant if $(x_0, y_0) \in D$, we have that the solution $x(T) := x(T; 0, x_0, y_0), y(T) := y(T; 0, x_0, y_0)$ belongs also to D . Considering that D is compact and simply connected, the Poincaré map has a fixed point, which of course is the initial condition of a T -periodic solution $\varphi(t)$ of system (5), and such solution verifies (4). \square

3. Asymptotically stability of periodic solutions

In this section we combine the bounds obtained in the proof for existence with the results in [19] in order to get a uniqueness and stability criterion.

THEOREM 2. *Under the condition of Theorem 1, assume moreover that g has a continuous derivative in its domain. Let us call*

$$m = \min_{x \in [l'_1, l'_2]} f(x), \quad M = \max_{x \in [l'_1, l'_2]} f(x)$$

and fix the constants

$$\beta = (M - m)/2, \quad \gamma = (M + m)/2, \quad \alpha = (\pi/T)^2 + \gamma^2/4.$$

If the following condition holds

$$0 < \max_{x \in [l'_1, l'_2]} g'(x) \leq \alpha - \beta(\gamma + \alpha^{1/2}), \tag{15}$$

then the periodic solution found in Theorem 1 is unique and asymptotically stable.

Proof. The uniqueness follows by using the argument of the proof of [19, Proposition 4.3]. Concerning the asymptotic stability, the solution found in Theorem 1 has index 1 because it comes from Brouwer's fixed point theorem, then we can apply [19, Proposition 1.2] (see also [19, Remark 1]). \square

4. Examples and comparison with related results

It is important to remark that the equation with weak singularities and nonlinear friction term has been scarcely explored until now. A recent reference is [18]. In order to compare the respective results, let us take the model equation

$$\ddot{x} + f(x)\dot{x} + x^\mu - \frac{1}{x^\delta} = a \cos(wt), \tag{16}$$

where $a, \mu, \delta, w > 0$. This corresponds to eq. (1) with

$$g(x) = x^\mu - \frac{1}{x^\delta} \text{ and } h(t) = a \cos(wt).$$

Of course $T = \frac{2\pi}{w}$. If (H2) holds, [18, Theorem 1.1] establishes the existence of a positive T -periodic solution under the condition

$$f_0 > \frac{\sqrt{T} \|h\|_2}{g^{-1}(-a)} = \frac{\pi\sqrt{2}a}{g^{-1}(-a)w}. \tag{17}$$

If compared with this condition, our main assumption (3) has the advantage that it does not depend on the frequency w . Therefore both assumptions are independent and it is easy to construct examples verifying (3) but not (17), just taking w small enough. For instance, fix $f(x) = 1, \mu = 1, \delta = \frac{1}{2}$, if we take $C = 1$ in Theorem 1, l'_1, l'_2 can be computed numerically giving $l'_1 = 0.340298, l'_2 = 1.73068$. As a consequence, $K_1 = 0.970549, K_2 = 0.589254$. Then, condition (3) reads

$$a \leq \frac{1}{2} \min\{K_1, K_2\} = 0.294627.$$

Under such condition, (16) has a T -periodic solution for all w . If $a = \frac{1}{4}$, a numerical computation shows that (17) does not hold if $w < 0.189215$. As a further example, if we take:

$$\mu = 2, \delta = \frac{1}{2}, f(x) \equiv 4, l'_1 = \frac{1}{2}, \text{ and } C = Z(l'_1),$$

then $l'_2 = 1.49107, K_1 = 0.698446, K_2 = 0.291053$, condition (3) reads

$$a \leq 2 \min\{K_1, K_2\} = 0.582107$$

and condition (15) reads

$$\max_{x \in [l'_1, l'_2]} g'(x) = 3.25676 < 4 + \frac{w^2}{4},$$

for any frequency $w > 0$. Therefore, if we choose $a = \frac{1}{2}$ we have the Theorems 1 and 2 guarantee the existence of a unique T -periodic solution which is asymptotically stable for all w , but a numerical computation shows that (17) does not hold if $w < 0.698132$.

Other interesting reference is [11]. In this paper, the authors study the equation with linear friction term ($f(x) \equiv c$ constant) by a combination of the method of upper and lower solutions and the ideas from [19]. In particular, for the equation

$$\ddot{x} + c\dot{x} - \frac{1}{x^\delta} = -1 + a \cos(\omega t), \quad (18)$$

with $a, c > 0$, Theorem 1.2 (see also Example 3.1) gives the existence of a unique T -periodic solution which is asymptotically stable under the condition

$$1 + a \leq \left(\frac{1}{4\delta}(w^2 + c^2) \right)^{\delta/(\delta+1)}. \quad (19)$$

Again, the condition depends explicitly on the frequency w , hence it is essentially independent from condition (3), being not difficult to derive explicit examples to illustrate this fact. We omit further details.

Acknowledgements. We are grateful to Prof. Rafael Ortega for pointing us the key reference [10].

REFERENCES

- [1] J. CHU AND P. J. TORRES, *Applications of Schauder's fixed point theorem to singular differential equations*, Bull. Lond. Math. Soc., **39**, 4 (2007), 653–660.
- [2] L. DERWIDUÉ, *Systèmes différentiels non linéaires ayant des solutions périodiques*, Acad. Roy. Belg. Bull. Cl. Sci., **49** (1963), 11–32.
- [3] R. FAURÉ, *Solutions périodiques d'équations différentielles et méthode de Leray-Schauder. (Cas des vibrations forcées)*, Ann. Inst. Fourier (Grenoble), **14** (1964), 195–204.
- [4] N. FORBAT AND A. HUAUX, *Détermination approchée et stabilité locale de la solution périodique d'une équation différentielle non linéaire*, Mém. Publ. Soc. Sci. Arts Lett. Hainaut, **76** (1962), 193–203.
- [5] A. HUAUX, *Sur l'existence d'une solution périodique de l'équation différentielle non linéaire $x'' + 0,2x' + x/(1-x) = (0,5)\cos \omega t$* , Bull. Cl. Sciences Acad. R. Belgique, **48**, 5 (1962), 494–504.
- [6] P. HABETS AND L. SANCHEZ, *Periodic solutions of some Liénard equations with singularities*, Proc. Amer. Math. Soc., **109**, 4 (1990), 1035–1044.
- [7] A. C. LAZER AND S. SOLIMINI, *On periodic solutions of nonlinear differential equations with singularities*, Proc. Amer. Math. Soc., **99**, 1 (1987), 109–114.
- [8] R. F. MARTINS, *Existence of periodic solutions for second-order differential equations with singularities and the strong force condition*, J. Math. Anal. Appl., **317**, 1 (2006), 1–13.
- [9] J. MAWHIN, *Topological degree and boundary value problems for nonlinear differential equations*, Topological methods for ordinary differential equations (Montecatini Terme, 1991), 74–142, Lecture Notes in Math., **1537**, Springer, Berlin, 1993.
- [10] M. NAGUMO, *On the periodic solution of an ordinary differential equation of second order*, Zenkoku Shijou Suugaku Danwakai, (1944), 54–61 (in Japanese), English translation in *Mitio Nagumo collected papers*, Springer-Verlag, 1993.
- [11] F.I. NJOKU AND P. OMARI, *Stability properties of periodic solutions of a Duffing equation in the presence of lower and upper solutions*, Appl. Math. Comput., **135**, (2003), 471–490.
- [12] I. RACHŮNKOVÁ, S. STANĚK AND M. TVRDÝ, *Singularities and Laplacians in boundary value problems for nonlinear ordinary differential equations*, Handbook of differential equations: ordinary differential equations. Vol. III, 607–722, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2006.

- [13] I. RACHŮNKOVÁ, S. STANĚK AND M. TVRDÝ, *Solvability of Nonlinear Singular Problems for Ordinary Differential Equations*, Contemporary Mathematics and Its Applications, **5**, Hindawi Publishing Corporation, New York, 2008.
- [14] I. RACHŮNKOVÁ, M. TVRDÝ AND I. VRKOČ, *Existence of nonnegative and nonpositive solutions for second-order periodic boundary-value problems*, J. Differential Equations, **176**, 2 (2001), 445–469.
- [15] P. J. TORRES, *Existence of one-signed periodic solutions of some second-order differential equations via a Krasnoselskii fixed point theorem*, J. Differential Equations, **190**, 2 (2003), 643–662.
- [16] P. J. TORRES, *Weak singularities may help periodic solutions to exist*, J. Differential Equations, **232**, 1 (2007), 277–284.
- [17] P. J. TORRES, *Existence and stability of periodic solutions for second-order semilinear differential equations with a singular nonlinearity*, Proc. Roy. Soc. Edinburgh Sect. A, **137**, 1 (2007), 195–201.
- [18] Z. ZHANG AND R. YUAN, *Existence of positive periodic solutions for the Liénard differential equations with weakly repulsive singularity*, Acta Appl. Math., **111**, 2 (2010), 171–178.
- [19] A. ZITAN AND R. ORTEGA, *Existence of asymptotically stable periodic solutions of a forced equation of Liénard type*, Nonlinear Anal., **22** (1994), 993–1003.

(Received January 13, 2011)

(Revised March 25, 2011)

Alexander Gutiérrez
Departamento de Matemáticas
Universidad Tecnológica de Pereira
Risaralda
Colombia
e-mail: alexquti@utp.edu.co

Pedro J. Torres
Departamento de Matemática Aplicada Universidad de Granada
18071 Granada
Spain
e-mail: ptorres@ugr.es