

Chaos in discrete structured population models

EDUARDO LIZ* AND ALFONSO RUIZ-HERRERA†

June 26, 2012

Abstract. We prove analytically the existence of chaotic dynamics in some classical discrete-time age-structured population models. Our approach allows us to estimate the sensitive dependence on the initial conditions, regions of initial data with chaotic behavior, and explicit ranges of parameters for which the considered models display chaos. These properties have important implications to evaluate the influence of a chaotic regime in the predictions based on mathematical models. We illustrate through particular examples how to apply our results.

Key Words: Chaotic dynamics; coin-tossing; structured population models; Clark equation.

AMS subject classifications (2010). 37D45, 37N25, 39A33, 92D25

1 Introduction

Understanding chaos is an important aim in many disciplines. May [24, 25] has already pointed out the possibility of simple deterministic models to display complex behavior, and the importance of chaos in ecology. The main implication of erratic fluctuations typical of a chaotic system is that long-time predictions are not possible; as emphasized by Hastings et al. [15], the characteristic of chaos that better represents this difficulty is a sensitive dependence on initial conditions. Besides this property, there are other important features in a chaotic system. Intuitively, and according to Smale's expository article [28], a chaotic phenomenon occurs if it is possible to reproduce, within the system and varying the initial conditions, all the possible outcomes of a coin-flipping experiment (see Definition 3.1). This notion has strong implications such as sensitive

*Departamento de Matemática Aplicada II, E. I. Telecomunicación, Universidad de Vigo, 36310 Vigo, Spain (eliz@dma.uvigo.es). Research partially supported by the Spanish Government and FEDER, grant MTM2010-14837.

†Departamento de Matemática Aplicada, Facultad de Ciencias, Universidad de Granada, 18071 Granada, Spain (alfonsoruiz@ugr.es). Research partially supported by the Spanish Government, project MTM2011-23652.

dependence on the initial conditions, and the existence of aperiodic orbits and infinitely many periodic points.

Although analytic proofs of chaos are now available for some one-dimensional maps usually employed in population dynamics (see, e.g, Thunberg's survey [30], and references therein), it is hard to find evidence of chaos in ecology. The main reason is the difficulty in manipulating and experimenting with ecological systems. A remarkable exception is the work made by Cushing et al. [7, 8, 10, 11]. They derived a structured population model, known as LPA, to study the growth of laboratory populations of flour beetles of the genus *Tribolium*, and their experiments confirmed chaotic behavior predicted by the model. Other age-structured population models in which chaos has been recently explored are Clark's equation [3, 16], and a class of density-dependent Leslie models [14, 19, 32, 33, 35].

The aim of this paper is twofold. On the one hand, we provide analytic proofs of existence of chaotic dynamics in some systems of difference equations, and we apply our results to the aforementioned structured population models. From a biological point of view, our strategy consists of introducing chaos in a specific age group, and then studying its influence in the full system; under a mathematical perspective, we derive criteria of chaotic dynamics in higher-dimensional systems using a generalization of some elementary notions of chaos in one dimension. On the other hand, an important purpose of this work is to interpret some natural properties of complex dynamics in ecological models. Specifically, we stress in the estimation of sensitive dependence on the initial conditions and in the location of regions of both the phase space and the set of parameters where the system displays chaos. These two properties have important practical implications to evaluate the influence of a chaotic regime in predictions based on mathematical models.

The organization of the paper is as follows. In Section 2, we review the specific population models in which we focus our study; in Section 3 we introduce the notion of chaotic dynamics that we use, and derive some results about chaos for continuous higher-dimensional maps. Sections 4 and 5 contain our main applications to prove chaos in the structured models introduced in Section 2; some computations are included in two appendices. Finally, in Section 6 we discuss the main conclusions of this paper.

2 Structured population models

Mathematical formulation of discrete models leads to consider difference equations or matrix models, which are nonlinear if changes in response to population density are taken into account. We focus our study on discrete-time age-structured models for a single population, that is to say, the model describes the distribution of individuals among the possible categories of important age differences. The monographs [4, 9, 31] are good sources both for theory and applications of structured population models.

Perhaps the simplest formulation that combines age structure and density-dependent recruitment has the form

$$x_n = \alpha x_{n-1} + (1 - \alpha)f(x_{n-k}), \quad (2.1)$$

which is commonly known as the Clark model [3, 5], and has been employed in fishery models (see [26] and its references). For $k = 1$, (2.1) is a first-order difference equation that provides a simple way to allow for certain survivorship of adults from one reproduction period to the next. This one-dimensional model was formulated by Clark [6]; see also [29, Section 9.2]. For recent work involving this equation, including chaotic dynamics, we refer the reader to [21, 23, 36].

For $k \geq 2$, we do not know any analytic proof of chaos in (2.1). Botsford [3] has already reported some numerical results suggesting chaotic behavior in (2.1) with the Ricker map $f(x) = xe^{r(1-x)}$, $r > 0$. For a more detailed study, including numerical computations of Lyapunov exponents for $k = 3$, we refer to [16].

Another simple structured model for which existence of chaos has been explored is a density-dependent Leslie population model with two age classes

$$\begin{cases} x_{n+1} = (a_1x_n + a_2y_n) \exp(-\lambda_1x_n - \lambda_2y_n) \\ y_{n+1} = bx_n \end{cases} \quad (2.2)$$

where $a_1, a_2, \lambda_1, \lambda_2$ are positive parameters, and $b \in]0, 1]$.

Guckenheimer et al. [14] considered the case $a_1 = a_2 = a$, $\lambda_1 = \lambda_2 = 0.1$, $b = 1$ to show numerically a route to chaos as a is increased. See also [32, 35] for a detailed study of stability and bifurcations. Recently, Ugarcovici and Weiss [33] proved the existence of a set of parameters for which system (2.2) admits a chaotic attractor.

As mentioned before, a study to predict the population dynamics of the flour beetle *Tribolium* published by Cushing et al. in a series of papers (see [11] and references therein) provides a structured population model that became a paradigm for chaos in ecology. The model distinguishes three stages (larvae, pupae and adults), and it is referred to as the LPA model:

$$\begin{cases} L_{n+1} = bA_n \exp(-c_{el}L_n - c_{ea}A_n) \\ P_{n+1} = (1 - \mu_l)L_n \\ A_{n+1} = P_n \exp(-c_{pa}A_n) + (1 - \mu_a)A_n \end{cases} \quad (2.3)$$

All coefficients are nonnegative, and $\mu_l < 1$, $\mu_a \leq 1$. See, e.g., [12] for a biological interpretation of the involved parameters and many properties of this system. A route to chaos numerically predicted by the LPA model (2.3) was confirmed by some laboratory experiments. However, as stated in [11], no proof of chaotic dynamics is available.

3 Background on chaotic dynamics

In this section we give the main tools and definitions that we use in this paper. We understand chaos in the sense of the following definition.

Definition 3.1. Consider (X, d) a metric space. We say that a continuous map $\psi : X \rightarrow X$ induces **chaotic dynamics on two symbols** if there exist two disjoint compact sets $\mathcal{K}_0, \mathcal{K}_1 \subset X$ such that, for each two-sided sequence $(s_i)_{i \in \mathbb{Z}} \in \{0, 1\}^{\mathbb{Z}}$,

there exists a corresponding sequence $(\omega_i)_{i \in \mathbb{Z}} \in (\mathcal{K}_0 \cup \mathcal{K}_1)^{\mathbb{Z}}$ such that

$$\omega_i \in \mathcal{K}_{s_i} \quad \text{and} \quad \omega_{i+1} = \psi(\omega_i) \quad \text{for all } i \in \mathbb{Z}, \quad (3.1)$$

and, whenever $(s_i)_{i \in \mathbb{Z}}$ is a k -periodic sequence (that is, $s_{i+k} = s_i, \forall i \in \mathbb{Z}$) for some $k \geq 1$, there exists a k -periodic sequence $(\omega_i)_{i \in \mathbb{Z}} \in (\mathcal{K}_0 \cup \mathcal{K}_1)^{\mathbb{Z}}$ satisfying (3.1).

Definition 3.1 guarantees natural properties of complex dynamics such as sensitive dependence on the initial conditions or the presence of an invariant set Λ being transitive and semi conjugate with the Bernoulli shift. See Theorem 2.2 in [27] for a list of properties of the map ψ that hold if the conditions of Definition 3.1 are satisfied. We note that a map that is chaotic according to Definition 3.1 is also chaotic in the sense of Block and Coppel and in the sense of *coin-tossing*; we refer the reader to [18] and [1, Remarks 3.2] for precise definitions and further comments. Moreover, our definition of chaos ensures the existence of periodic points of any period $n \in \mathbb{N}$, in contrast with other definitions (see, e.g., [1, 2, 18]).

Our primary purpose is to obtain some criteria to detect chaos analytically. To this end, we use the method developed by Zgliczyński and Gidea in [37] (see also [38]). Definitions 3.2 and 3.3 below correspond to Definitions 2 and 3 in [37]. We employ the usual maximum norm in \mathbb{R}^n ,

$$\|(x_1, x_2, \dots, x_n)\| = \max\{|x_i| : i = 1, 2, \dots, n\},$$

and use the notation $J_n = [-1, 1]^n$ for the closed ball of radius 1 centered at $0 \in \mathbb{R}^n$.

Definition 3.2. *An h -set is a quadruple consisting of*

- a compact subset N of \mathbb{R}^n ,
- a pair of numbers $u = u(N), s = s(N) \in \{0, 1, 2, \dots\}$, with $u + s = n$,
- a homeomorphism $c_N : \mathbb{R}^n \rightarrow \mathbb{R}^n$, such that $c_N(N) = J_n$.

In this setting, we employ the notation

$$N_c^- = \partial J_u \times J_s,$$

$$N_c^+ = J_u \times \partial J_s.$$

As mentioned in [37], the numbers $u(N)$ and $s(N)$ stand for the dimensions of nominally unstable and stable directions, respectively. Notice that if $u(N) = 0$ then $N_c^- = \emptyset$, and if $s(N) = 0$ then $N_c^+ = \emptyset$.

Definition 3.3. *Assume that N, M are h -sets, such that $u(N) = u(M) = u$ and $s(N) = s(M) = s$. Let $f : N \rightarrow \mathbb{R}^n$ be a continuous map, and define $f_c = c_M \circ f \circ c_N^{-1} : J_n \rightarrow \mathbb{R}^n$. We say that N f -covers M , and write it as*

$$N \xrightarrow{f} M,$$

if the following conditions are satisfied:

- 1 *There exists a continuous homotopy $H : [0, 1] \times J_n \rightarrow \mathbb{R}^n$, such that the following conditions hold true:*

$$H_0 = f_c,$$

$$H([0, 1], N_c^-) \cap J_n = \emptyset,$$

$$H([0, 1], J_n) \cap M_c^+ = \emptyset.$$

- 2 *There exists a linear map $A : \mathbb{R}^u \rightarrow \mathbb{R}^u$, such that $H_1(p, q) = (Ap, 0)$ for $p \in J_u$ and $q \in J_s$, and $A(\partial J_u) \subset \mathbb{R}^u \setminus J_u$.*

Next we link these concepts with existence of chaotic dynamics.

Theorem 3.1. *Let $F : \mathcal{D} \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous map and assume that there exist two disjoint h-sets N_0 and N_1 such that*

$$N_i \xrightarrow{F} N_j$$

for all $i, j = 0, 1$. Then F induces chaotic dynamics on two symbols (with compact sets $\mathcal{K}_0 = N_0$ and $\mathcal{K}_1 = N_1$).

Proof. Let us first take a sequence $s = (s_i) \in \{0, 1\}^{\mathbb{Z}}$ so that $s_{i+k} = s_i$ for all $i \in \mathbb{Z}$. In this case, it follows from Theorem 4 in [37] that there exists $x \in N_0 \cup N_1$ so that

$$F^k(x) = x, \text{ and } F^i(x) \in N_{s_i}, \text{ for all } i \in \mathbb{Z}.$$

Now we prove the assertion of the theorem. Fix an arbitrary sequence $s = (s_j)_{j \in \mathbb{Z}}$, and, for each $i \in \mathbb{N}$, define the compact set

$$\Gamma_i = \{\omega \in N_{s_0} : F^j(\omega) \in N_{s_j} \text{ for all } 1 \leq j \leq i\}.$$

Clearly $\Gamma_{i+1} \subset \Gamma_i$ and, by the previous step, $\Gamma_i \neq \emptyset$, for all $i \in \mathbb{N}$. Indeed, it is sufficient to consider the $(i+1)$ -periodic sequence that matches s for $j = 0, 1, \dots, i$. Hence,

$$\bigcap_{i=1}^{\infty} \Gamma_i \neq \emptyset.$$

To conclude the proof of the theorem, we notice that a standard diagonal argument enables us to extend the result to bi-infinite sequences; see [17, Theorem 2.2]. \square

The following result provides us with an elementary and effective method to estimate the sensitive dependence in a chaotic regime when the conditions of Theorem 3.1 hold.

Proposition 3.1. *Take N_0, N_1 , and $F : \mathcal{D} \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ as in the statement of Theorem 3.1, and denote $d = \text{dist}(N_0, N_1) > 0$. For $\varepsilon > 0$, we define*

$$S_\varepsilon = \max\{n \in \mathbb{N} : N_0 \text{ contains } n \text{ disjoint balls of diameter } \varepsilon\},$$

and

$$N^* = 1 + \left\lceil \frac{\ln S_\varepsilon}{\ln 2} \right\rceil,$$

where, for $x \in \mathbb{R}$, $\lceil x \rceil$ denotes the ceiling of x , that is, the smallest integer not less than x .

Then, there are two points x_0, y_0 satisfying that

- $x_0, y_0 \in N_0$,
- $\|x_0 - y_0\| < \varepsilon$,
- $\max_{0 \leq j \leq N^*} \{\|F^j(x_0) - F^j(y_0)\|\} > d$.

Proof. Fix $\varepsilon > 0$. Given $\tau = (\tau_1, \dots, \tau_{N^*})$ with $\tau_i \in \{0, 1\}$ we define

$$\Gamma_\tau = \{x \in N_0 : F^j(x) \in N_{\tau_j} \text{ for all } 1 \leq j \leq N^*\}.$$

In this way, we can construct 2^{N^*} disjoint nonempty compact sets contained in N_0 . Since, by definition, $S_\varepsilon < 2^{N^*}$, there are two different N^* -tuples, τ', τ'' so that

$$\text{dist}(\Gamma_{\tau'}, \Gamma_{\tau''}) < \varepsilon.$$

Hence, we can choose two points $x_0 \in \Gamma_{\tau'}$ and $y_0 \in \Gamma_{\tau''}$ so that $\|x_0 - y_0\| < \varepsilon$.

Since $\tau' \neq \tau''$, there exists at least one index $j_0 \leq N^*$ such that $\tau'_{j_0} \neq \tau''_{j_0}$. By the definition of $\Gamma_{\tau'}$ and $\Gamma_{\tau''}$, it follows that

$$F^{j_0}(x_0) \in N_{\tau'_{j_0}}, \quad F^{j_0}(y_0) \in N_{\tau''_{j_0}},$$

and therefore $\|F^{j_0}(x_0) - F^{j_0}(y_0)\| \geq \text{dist}(N_0, N_1) = d$, as we wanted to prove. \square

4 Chaotic dynamics in models (2.1) and (2.3)

In this section we illustrate how to apply the previous results to systems (2.1) and (2.3). For these models, we obtain analytically chaotic regimes where the map of the system only has unstable directions in the sense of Definition 3.2. To this aim, we need to introduce the following notion of turbulence, which is more restrictive than the usual one [2, Chapter II].

Definition 4.1. *Let I be a real interval, and $g : I \rightarrow I$ a continuous map. We say that g is δ -strictly turbulent if there exist four constants $\beta_0 < \beta_1 < \gamma_0 < \gamma_1$, and $\delta > 0$ so that*

$$\begin{aligned} g(\beta_0) &< \beta_0 - \delta < \gamma_1 + \delta < g(\beta_1), \\ g(\gamma_1) &< \beta_0 - \delta < \gamma_1 + \delta < g(\gamma_0). \end{aligned}$$

We first consider the Clark model (2.1). There is a one-to-one correspondence between the solutions of equation (2.1) and the orbits of the map $F_\alpha : \mathbb{R}^k \rightarrow \mathbb{R}^k$ defined by

$$F_\alpha(x_1, \dots, x_k) = (x_2, \dots, x_k, \alpha x_k + (1 - \alpha)f(x_1)).$$

For this map we get the following result.

Theorem 4.1. *Assume that f is δ -strictly turbulent with parameters $\beta_0 < \beta_1 < \gamma_0 < \gamma_1$. Then there exists $\alpha_0 > 0$ so that for all $0 \leq \alpha \leq \alpha_0$, F_α^k induces chaotic dynamics on two symbols relative to $N_0 = [\beta_0, \beta_1]^k$ and $N_1 = [\gamma_0, \gamma_1]^k$.*

Proof. It follows from the expression of F_α that, for $\alpha = 0$,

$$F_0^k(x_1, \dots, x_k) = (f(x_1), \dots, f(x_k)).$$

Using the continuity of F_α with respect to α , we obtain that there is a constant $\alpha_0 > 0$ so that for all $i = 1, \dots, k$, $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k) \in [\beta_0, \beta_1]^{k-1}$, and $\alpha \leq \alpha_0$,

$$|(F_\alpha^k)_i(x_1, \dots, x_{i-1}, \beta_0, x_{i+1}, \dots, x_k) - f(\beta_0)| < \delta, \quad (4.1)$$

$$|(F_\alpha^k)_i(x_1, \dots, x_{i-1}, \beta_1, x_{i+1}, \dots, x_k) - f(\beta_1)| < \delta. \quad (4.2)$$

Analogously, for all $i = 1, \dots, k$ and $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k) \in [\gamma_0, \gamma_1]^{k-1}$,

$$|(F_\alpha^k)_i(x_1, \dots, x_{i-1}, \gamma_0, x_{i+1}, \dots, x_k) - f(\gamma_0)| < \delta, \quad (4.3)$$

$$|(F_\alpha^k)_i(x_1, \dots, x_{i-1}, \gamma_1, x_{i+1}, \dots, x_k) - f(\gamma_1)| < \delta. \quad (4.4)$$

Next, let us consider the translations t_v, t_w according to the vectors v, w given by

$$v = \left(-\frac{\beta_0 + \beta_1}{2}, \dots, -\frac{\beta_0 + \beta_1}{2} \right), \quad w = \left(-\frac{\gamma_0 + \gamma_1}{2}, \dots, -\frac{\gamma_0 + \gamma_1}{2} \right),$$

respectively, and the maps h_0, h_1 defined by

$$h_0(x_1, \dots, x_k) = \frac{2}{\beta_1 - \beta_0}(x_1, \dots, x_k), \quad h_1(x_1, \dots, x_k) = \frac{2}{\gamma_1 - \gamma_0}(x_1, \dots, x_k).$$

We claim that the h-cubes N_0, N_1 with

- $u(N_0) = u(N_1) = k$ and $s(N_0) = s(N_1) = 0$,
- $c_{N_0} = h_0 \circ t_v, c_{N_1} = h_1 \circ t_w$,

satisfy the covering relations

$$N_i \xrightarrow{F_\alpha^k} N_j,$$

for all $i, j = 0, 1$. Note that $N_c^+ = \emptyset$ and $N_c^- = \partial J_k$, for $N = N_i, i = 0, 1$.

We first prove the relation

$$N_0 \xrightarrow{F_\alpha^k} N_0. \quad (4.5)$$

To this end, we take the linear map $A(x) = 2x$, and define the homotopy $H : [0, 1] \times J_k \rightarrow \mathbb{R}^k$ by

$$H(t, x) = tA(x) + (1 - t)(c_{N_0} \circ F_\alpha^k \circ c_{N_0}^{-1})(x).$$

Clearly, $H(0, \cdot) = c_{N_0} \circ F_\alpha^k \circ c_{N_0}^{-1}$ and $H(1, \cdot) = A$. Next we prove that

$$H(t, \partial J_k) \cap J_k = \emptyset.$$

Indeed, take a point $x \in \partial J_k$. For this point, we can ensure that there is an index $i \in \{1, \dots, k\}$ so that x_i is either equal to -1 or to 1 . Assume that we are in the first case (the proof in the other case is completely analogous). By the definition of c_{N_0} , we deduce that

$$c_{N_0}^{-1}(x) = (y_1, \dots, y_{i-1}, \beta_0, y_{i+1}, \dots, y_k),$$

with $(y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_k) \in [\beta_0, \beta_1]^{k-1}$. Now, using condition (4.1), it follows that

$$|(F_\alpha^k)_i(y_1, \dots, y_{i-1}, \beta_0, y_{i+1}, \dots, y_k) - f(\beta_0)| < \delta.$$

Therefore, as f is δ -strictly turbulent, we obtain that

$$(F_\alpha^k)_i(y_1, \dots, y_{i-1}, \beta_0, y_{i+1}, \dots, y_k) < \beta_0.$$

Finally, by the definition of c_{N_0} we arrive at

$$(c_{N_0} \circ F_\alpha^k)_i(y_1, \dots, y_{i-1}, \beta_0, y_{i+1}, \dots, y_k) < -1.$$

Bringing all the information together, and using that $(Ax)_i = 2x_i = -2$, it is clear that

$$t(Ax)_i + (1 - t)(c_{N_0} \circ F_\alpha^k \circ c_{N_0}^{-1}(x))_i < -1,$$

for all $t \in [0, 1]$, and therefore $H(t, x) \notin J_k$. The proof of (4.5) is now complete. The same reasoning applies to check the remainder covering relations. We just note that for the relations

$$N_1 \xrightarrow{F_\alpha^k} N_i$$

with $i = 0, 1$, the choice for the linear map is $A(x) = -2x$. □

Remark 4.1. *If some iteration f^m is δ -strictly turbulent, then the conclusion of Theorem 4.1 holds replacing F_α^k by F_α^{km} .*

Remark 4.2. *The requirement of an upper bound in the range of α imposed in Theorem 4.1 is necessary. For values of α close to 1, equation (2.1) has a globally stable positive equilibrium (see [13, 20] and their references). Actually, for $k = 1$, α is sometimes used as a control parameter so that increasing its value becomes an effective tool for chaos control [22].*

To ensure the existence of chaotic dynamics in (2.1) for a particular choice of α when f is δ -strictly turbulent, we have to verify conditions (4.1)-(4.4). We illustrate this fact with an example.

Example 4.1. Consider the system

$$x_n = \alpha x_{n-1} + (1 - \alpha)f(x_{n-2}), \quad (4.6)$$

where $f(x) = x \exp(4 - x)$. A direct computation allows us to check that f^2 is 2.4-strictly turbulent with parameters $3 < 5.75 < 6.25 < 11$. In Appendix 1, we show that conditions (4.1)-(4.4) hold for f^2 with $\alpha_0 = 0.001$.

It is worth pointing out some biological implications of our results for equation (4.6). Specifically, there are two disjoint regions, namely

$$N_0 = [3, 5.75]^2, \quad N_1 = [6.25, 11]^2,$$

with the ‘‘coin-tossing’’ property for F_α^4 . For instance, if we take the bi-infinite sequence

$$(\dots, 1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 1, \dots)$$

(assume that the first 1 corresponds to the index 0 in the sequence), we can choose a point x so that

$$x \in N_1, \quad F_\alpha^4(x) \in N_0, \quad F_\alpha^8(x) \in N_1, \quad F_\alpha^{12}(x) \in N_0, \quad F_\alpha^{16}(x) \in N_0, \quad F_\alpha^{20}(x) \in N_1, \dots$$

Moreover, if the sequence is periodic, we can take a periodic point following the ‘‘itinerary’’ of the sequence. As a direct consequence of Proposition 3.1, we can also estimate the sensitive dependence. Indeed, for $\varepsilon > 0$, clearly

$$S_\varepsilon \leq \left\lceil \frac{(5.75 - 3)^2}{\varepsilon^2} \right\rceil := B$$

Hence, there are two points $x_0, y_0 \in [3, 5.75]^2$ such that $\|x_0 - y_0\| < \varepsilon$ and, for some number $j \in \{1, \dots, N^*\}$, we have that $F^{4j}(x_0) \in [3, 5.75]^2$ and $F^{4j}(y_0) \in [6.25, 11]^2$. Therefore,

$$\|F^{4j}(x_0) - F^{4j}(y_0)\| \geq 0.5.$$

We emphasize that, for any given value of $\varepsilon > 0$, we can compute an explicit bound $\lceil \frac{\ln B}{\ln 2} \rceil + 1$ for N^* .

Next we study the LPA model (2.3), whose solutions are the orbits of the map $G : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by

$$G(L, P, A) = (bA \exp(-c_{el}L - c_{ea}A), (1 - \mu_l)L, P \exp(-c_{pa}A) + (1 - \mu_a)A). \quad (4.7)$$

Clearly, for $c_{el} = \mu_l = c_{pa} = 0$ and $\mu_a = 1$, system (2.3) becomes Clark’s model with $\alpha = 0$, $k = 3$ and $f(x) = bx e^{-c_{ea}x}$. Thus, the proof of Theorem 4.1 can be easily adapted to deal with (2.3). Since there are values of b and c_{ea} so that f^2 is δ -strictly turbulent, the following result establishes chaos for the LPA model.

Theorem 4.2. *Assume that f^2 is δ -strictly turbulent with parameters $\beta_0 < \beta_1 < \gamma_0 < \gamma_1$, and $\delta > 0$. There exists $\sigma > 0$ so that if $0 \leq c_{el}, \mu_l, c_{pa} \leq \sigma$ and $1 - \sigma \leq \mu_a \leq 1$, then G^6 induces chaotic dynamics on two symbols relative to $N_0 = [\beta_0, \beta_1]^3$ and $N_1 = [\gamma_0, \gamma_1]^3$, where G is the map defined in (4.7).*

We observe that, reasoning in a similar way as in the previous example, it is possible to estimate properties of sensitive dependence on the initial conditions for particular cases of system (2.3).

To the best of the authors' knowledge, Theorem 4.2 is the first analytic result of chaotic dynamics for the LPA model in the interior of the nonnegative cone \mathbb{R}_+^3 . It is worth mentioning that in [10, Theorem 2], Cushing proves analytically the existence of fully synchronous orbits with chaotic behavior when $\mu_a = 1$.

5 Chaotic dynamics in system (2.2)

In this section we apply the results of Section 3 to the nonlinear Leslie population model (2.2). In contrast to the previous section, in this case we find a chaotic regime where the map of the system has one stable direction and other one unstable (in the sense of Definition 3.2). Associated to (2.2), we define the map in \mathbb{R}^2

$$F(x, y) = ((a_1x + a_2y) \exp(-\lambda_1x - \lambda_2y), bx),$$

where all the parameters are strictly positive, and $b \in]0, 1]$. Our main result in this section is the following.

Theorem 5.1. *Assume that $f(x) = a_1x \exp(-\lambda_1x)$ satisfies that f^2 is δ -strictly turbulent with parameters $\alpha_0 < \alpha_1 < \beta_0 < \beta_1$ and $\delta > 0$. Suppose that there is $r > 0$ so that the following inequalities are fulfilled:*

$$\frac{b}{e} \left(\frac{a_1}{\lambda_1} + \frac{a_2}{\lambda_2} \right) < r \quad (5.1)$$

$$\frac{-a_1}{e} (e^{-\lambda_2 r} - 1) < \ln \left(\frac{\alpha_0 - a_1 a_2 r - a_2 b \beta_1}{\alpha_0 - \delta} \right) \quad (5.2)$$

$$\lambda_2 r + \lambda_2 b \beta_1 + \lambda_1 a_2 r < \ln \left(\frac{\beta_1 + \delta}{\beta_1} \right). \quad (5.3)$$

Then F^2 induces chaotic dynamics on two symbols relative to $N_0 = [\alpha_0, \alpha_1] \times [0, r]$ and $N_1 = [\beta_0, \beta_1] \times [0, r]$.

Proof. It is easy to check that N_0 and N_1 are h -sets, with

- $u(N_0) = u(N_1) = 1$ (x -direction) and $s(N_0) = s(N_1) = 1$ (y -direction),
- $c_{N_0} = h_0 \circ t_v$ and $c_{N_1} = h_1 \circ t_w$,

where t_v and t_w are the translations according to the vectors

$$v = \left(\frac{-(\alpha_0 + \alpha_1)}{2}, \frac{-r}{2} \right), \quad w = \left(\frac{-(\beta_0 + \beta_1)}{2}, \frac{-r}{2} \right),$$

respectively, and

$$h_0(x, y) = \left(\frac{2x}{\alpha_1 - \alpha_0}, \frac{2y}{r} \right), \quad h_1(x, y) = \left(\frac{2x}{\beta_1 - \beta_0}, \frac{2y}{r} \right).$$

We have to demonstrate that

$$N_i \xrightarrow{F^2} N_j,$$

for $i, j = 0, 1$. We give the proof only for the case $i = j = 0$. Indeed, consider the homotopy

$$H(t, (x, y)) = t(c_{N_0} \circ F^2 \circ c_{N_0}^{-1})(x, y) + (1-t)A(x, y),$$

where $A(x, y) = (2x, 0)$.

Define $f(x) = a_1 x e^{-\lambda_1 x}$. After some elementary computations (see Appendix 2), we arrive at the inequalities

$$\begin{aligned} F_1^2(x, y) &\leq f^2(x)(\exp(-a_1/e))^{\exp(-\lambda_2 y)-1} + a_1 a_2 y + a_2 b x \\ F_1^2(x, y) &\geq f^2(x) \exp(-\lambda_2 b x) \exp(-\lambda_1 a_2 y) \exp(-\lambda_2 y) \\ F_2^2(x, y) &\leq b \left(\frac{a_1}{\lambda_1 e} + \frac{a_2}{\lambda_2 e} \right). \end{aligned}$$

Now, since $b > 0$, we deduce from (5.1) that, for all $(x, y) \in N_0$,

$$0 < F_2^2(x, y) < r. \quad (5.4)$$

Analogously, as $f^2(\alpha_0) < \alpha_0 - \delta$, we obtain that

$$F_1^2(\alpha_0, y) \leq (\alpha_0 - \delta)(\exp(-a_1/e))^{\exp(-\lambda_2 r)-1} + a_1 a_2 r + a_2 b \alpha_0, \quad (5.5)$$

for all $y \in [0, r]$. By using (5.2) together with the inequality $\alpha_0 < \beta_1$, we get from expression (5.5) that, for all $y \in [0, r]$,

$$F_1^2(\alpha_0, y) < \alpha_0. \quad (5.6)$$

Reasoning in the same way with condition (5.3), it may be concluded that

$$F_1^2(\alpha_1, y) > \alpha_1, \quad (5.7)$$

for all $y \in [0, r]$.

From inequalities (5.4), (5.6), and (5.7), it follows that

$$\begin{aligned} c_{N_0} \circ F^2 \circ c_{N_0}^{-1}(\{-1\} \times [-1, 1]) &\subset \{(x, y) : x < -1\} \\ c_{N_0} \circ F^2 \circ c_{N_0}^{-1}(\{1\} \times [-1, 1]) &\subset \{(x, y) : x > 1\} \\ c_{N_0} \circ F^2 \circ c_{N_0}^{-1}([-1, 1]^2) &\subset \{(x, y) : -1 < y < 1\}. \end{aligned}$$

These properties, together with the expression of A , lead to the desired conclusion

$$\begin{aligned} H([0, 1], \{-1, 1\} \times [-1, 1]) \cap [-1, 1]^2 &= \emptyset, \\ H([0, 1], [-1, 1]^2) \cap ([-1, 1] \times \{-1, 1\}) &= \emptyset. \end{aligned}$$

The same steps lead to the covering relation

$$N_0 \xrightarrow{F^2} N_1.$$

For the relations

$$N_1 \xrightarrow{F^2} N_i,$$

with $i = 0, 1$, the most significant change consists of taking the linear map $A(x, y) = (-2x, 0)$. \square

Now we apply Theorem 5.1 in a particular example. Take $f(x) = xe^{4-x}$ ($a_1 = e^4$, $\lambda_1 = 1$). As claimed in Section 4, f^2 is 2.4-strictly turbulent with parameters $3 < 5.75 < 6.25 < 11$. Straightforward computations show that conditions (5.1)-(5.3) hold for $r = 0.065$, $b = 0.003$, $a_2 = 0.1$, and $\lambda_2 = 1$.

In general, to use Theorem 5.1, the parameters b and r must be close to zero. Nevertheless, for values of b near 1, system (2.2) also displays chaotic dynamics; indeed, for $a_1 = \lambda_1 = 0$ and $b = 1$, we recover again Clark's model (2.1) with $\alpha = 0$. Consequently, arguing as in the proof of Theorem 4.1, we are able to guarantee the existence of chaos for (2.2) when (a_1, λ_1, b) belongs to a neighborhood of $(0, 0, 1)$. It is important to observe that, from a mathematical point of view, the chaotic regimes are different because in Theorem 5.1 we have one stable direction and one unstable direction, while in Theorem 4.1 the two directions are unstable.

6 Discussion

We have proved analytic criteria of chaotic dynamics for some well-known structured population models, and we have given further insight in such relevant features of chaos as the sensitive dependence on initial conditions. We list the main advantages of our results:

- **Explicit ranges of parameters where models (2.1), (2.2), and (2.3) exhibit chaotic behavior:** A key property from a biological point of view is the robustness of our results under small continuous perturbations. This aspect is especially important in modeling, because the involved coefficients are usually approximations derived from experimental data, so small errors should be taken into account.
- **Regions of initial data with chaotic behavior:** We find out explicit regions of the phase space, namely N_0 and N_1 , where chaotic behavior occurs. Determining

these regions is useful to analyze the influence of the chaotic regime on the system since, in some situations, a chaotic attractor may coexist with other dynamics, such as a locally stable periodic point.

- **Estimates of the sensitive dependence:** The strategy followed in our proofs allows us to get quantitative results about sensitive dependence on initial conditions in an easy way; see Proposition 3.1 and Example 4.1.

In the literature there are several papers dealing with the existence of chaotic dynamics for structured population models. However, most of them generally rely on numerical evidences obtained from computer simulations, in particular, on the computation of the dominant Lyapunov exponent (see, for instance, [16] for Clark's equation, and [11] for the LPA model).

For the nonlinear Leslie model (2.2), our results complement Theorem 1.2 in [33], which, based on the results of [34], ensures the existence of a chaotic attractor for a set of parameters with positive Lebesgue measure; however, compared with our approach, determining such a set can be much more involved. We note that our results only provide criteria for the existence of invariant sets with chaotic dynamics, not attractors. In particular, this complex behavior might not be observed in numerical simulations of the models. We point out that Theorem 1.2 in [33] establishes stronger attraction and stochastic properties.

Appendix 1

This appendix contains some computations for Example 4.1.

For $F_\alpha(x_1, x_2) = (x_2, \alpha x_2 + (1 - \alpha)f(x_1))$, the expression of $F_\alpha^4 = ((F_\alpha^4)_1, (F_\alpha^4)_2)$ is given by

$$\begin{aligned} (F_\alpha^4)_1(x_1, x_2) &= \alpha^3 x_2 + \alpha^2(1 - \alpha)f(x_1) + \alpha(1 - \alpha)f(x_2) \\ &\quad + (1 - \alpha)f(\alpha x_2 + (1 - \alpha)f(x_1)) \\ (F_\alpha^4)_2(x_1, x_2) &= \alpha^4 x_2 + \alpha^3(1 - \alpha)f(x_1) + \alpha^2(1 - \alpha)f(x_2) \\ &\quad + \alpha(1 - \alpha)f(\alpha x_2 + (1 - \alpha)f(x_1)) \\ &\quad + (1 - \alpha)f(\alpha^2 x_2 + \alpha(1 - \alpha)f(x_1) + (1 - \alpha)f(x_2)). \end{aligned}$$

Next we notice that, in $[0, +\infty[$, function $f(x) = x \exp(4 - x)$ is bounded by $\exp(3)$ and it is Lipschitz-continuous with Lipschitz-constant $\exp(4)$. Using these properties in a direct way, we obtain that, for all $x_2 \in [3, 5.75] \cup [6.25, 11]$,

$$\begin{aligned} |(F_\alpha^4)(x_1, x_2)_1 - f^2(x_1)| &\leq 11\alpha^3 + \exp(3)\alpha^2(1 - \alpha) + \exp(3)\alpha(1 - \alpha) \\ &\quad + \exp(4)(11\alpha + \exp(3)\alpha) + \alpha \exp(3) := \varphi(\alpha), \end{aligned}$$

and, for all $x_1 \in [3, 5.75] \cup [6.25, 11]$,

$$\begin{aligned} |(F_\alpha^4)(x_1, x_2))_2 - f^2(x_2)| \leq & 11\alpha^4 + \alpha^3(1 - \alpha) \exp(3) + \alpha^2(1 - \alpha) \exp(3) + \alpha(1 - \alpha) \exp(3) \\ & + 11 \exp(4)\alpha^2 + \exp(7)\alpha(1 - \alpha) + \exp(7)\alpha + \exp(3)\alpha := \psi(\alpha). \end{aligned}$$

Now, since φ and ψ are polynomials, it is elementary to check that $\varphi(\alpha) < 2.4$ and $\psi(\alpha) < 2.4$ for all $\alpha < \alpha_0 = 0.001$ (see Figure 1).




Figure 1: Representation of functions $\varphi(\alpha)$ and $\psi(\alpha)$ for $0 \leq \alpha \leq 0.002$.

Appendix 2

In this appendix, we provide some expressions and preliminary calculations for the proof of Theorem 5.1.

Consider the map

$$F(x, y) = ((a_1x + a_2y) \exp(-\lambda_1x - \lambda_2y), bx).$$

The first component of F^2 has the following expression:

$$\begin{aligned} F_1^2(x, y) = & e^{-\lambda_1 a_1 x} e^{-\lambda_1 a_1 x \exp(-\lambda_1 x - \lambda_2 y)} e^{-\lambda_2 b x} e^{-\lambda_1 a_2 y \exp(-\lambda_1 x - \lambda_2 y)} [a_1^2 x e^{-\lambda_1 x - \lambda_2 y} \\ & + a_1 a_2 y e^{-\lambda_1 x - \lambda_2 y} + a_2 b x]. \end{aligned}$$

We easily deduce that, for $x \geq 0$ and $y \geq 0$,

$$F_1^2(x, y) \leq a_1^2 x e^{-\lambda_1 x} e^{-\lambda_1 a_1 x \exp(-\lambda_1 x)} (e^{-\lambda_1 a_1 x \exp(-\lambda_1 x)})^{e^{-\lambda_2 y} - 1} + a_1 a_2 y + a_2 b x.$$

Now, using that $f(x) = a_1 x e^{-\lambda_1 x} \leq \frac{a_1}{\lambda_1 e}$, we arrive at

$$\begin{aligned} F_1^2(x, y) &\leq a_1^2 x e^{-\lambda_1 x} e^{-\lambda_1 a_1 x \exp(-\lambda_1 x)} \left(e^{\frac{-a_1}{e}} \right)^{e^{-\lambda_2 y} - 1} + a_1 a_2 y + a_2 b x \\ &= f^2(x) \left(e^{\frac{-a_1}{e}} \right)^{e^{-\lambda_2 y} - 1} + a_1 a_2 y + a_2 b x. \end{aligned}$$

On the other hand,

$$\begin{aligned} F_1^2(x, y) &\geq e^{-\lambda_1 a_1 x \exp(-\lambda_1 x - \lambda_2 y)} e^{-\lambda_2 b x} e^{-\lambda_1 a_2 y \exp(-\lambda_1 x - \lambda_2 y)} a_1^2 x e^{-\lambda_1 x - \lambda_2 y} \\ &\geq a_1^2 x e^{-\lambda_1 x} e^{-\lambda_1 a_1 x \exp(-\lambda_1 x)} e^{-\lambda_2 b x} e^{-\lambda_1 a_2 y} e^{-\lambda_2 y} \\ &= f^2(x) e^{-\lambda_2 b x} e^{-\lambda_1 a_2 y} e^{-\lambda_2 y}. \end{aligned}$$

For the second component, using that $f(x) = a_1 x e^{-\lambda_1 x} \leq \frac{a_1}{\lambda_1 e}$, we get

$$F_2^2(x, y) = b(a_1 x + a_2 y) e^{-\lambda_1 x - \lambda_2 y} \leq b \left(\frac{a_1}{\lambda_1 e} + \frac{a_2}{\lambda_2 e} \right).$$

Acknowledgements

We thank an anonymous referee for his/her useful comments. The second author thanks the Departamento de Matemática Aplicada II of the University of Vigo (Spain) for its kind hospitality.

References

- [1] B. AULBACH AND B. KIENINGER, *On three definitions of chaos*, *Nonlinear Dyn. Syst. Theory*, 1 (2001), pp. 23–37.
- [2] L.S. BLOCK AND W.A. COPPEL, *Dynamics in one dimension*, *Lectures Notes in Mathematics*, vol. 1513, Springer, Berlin.
- [3] L. W. BOTSFORD, *Further analysis of Clark's delayed recruitment model*, *Bull. Math. Biol.*, 54 (1992), pp. 275–293.
- [4] H. CASWELL, *Matrix Population Models: Construction, Analysis and Interpretation*, (2nd Ed.), Sinauer Associates, Sunderland, Massachusetts, 2001.
- [5] C. W. CLARK, *A delayed recruitment model of population dynamics with an application to baleen whale populations*, *J. Math. Biol.*, 3 (1976), pp. 381–391.
- [6] C. W. CLARK, *Mathematical Bioeconomics: The Optimal Management of Renewable Resources* (2nd Ed.), John Wiley & Sons, Hoboken, New Jersey, 1990.
- [7] R. F. COSTANTINO, J. M. CUSHING, B. DENNIS, AND R. A. DESHARNAIS, *Experimentally induced transitions in the dynamic behaviour of insect populations*, *Nature*, 375 (1995), pp. 227–230.

- [8] R. F. COSTANTINO, R. A. DESHARNAIS, J. M. CUSHING, AND B. DENNIS, *Chaotic dynamics in an insect population*, Science 275 (1997), pp. 389–391.
- [9] J. M. CUSHING, *An Introduction to Structured Population Dynamics*, CBMS-NSF Regional Conference Series in Applied Mathematics, 71, SIAM, Philadelphia, PA, 1998.
- [10] J. M. CUSHING, *Cycle chains and the LPA model*, J. Difference Equ. Appl., 9 (2003), pp. 655–670.
- [11] J. M. CUSHING, R. F. COSTANTINO, B. DENNIS, R. A. DESHARNAIS, AND S. M. HENSON, *Chaos in Ecology: Experimental Nonlinear Dynamics*, Theoretical Ecology Series, Academic Press, New York, 2003.
- [12] J. M. CUSHING, *The LPA model*, Fields Inst. Commun., 42 (2004), pp. 29–55.
- [13] H. A. EL-MORSHEDEY AND E. LIZ, *Globally attracting fixed points in higher order discrete population models*, J. Math. Biol., 53 (2006), pp. 365–384.
- [14] J. GUCKENHEIMER, G. OSTER, AND A. IPAKTCHI, *The dynamics of density dependent population models*, J. Math. Biol., 4 (1977), pp. 101–147.
- [15] A. HASTINGS, C. L. HOM, S. P. ELLNER, P. TURCHIN, AND H. C. J. GODFRAY, *Chaos in ecology: is mother nature a strange attractor?*, Ann. Rev. Ecol. Syst., 24 (1993), pp. 1–33.
- [16] K. HIGGINS, A. HASTINGS, AND L.W. BOTSFORD, *Density dependence and age structure: Nonlinear dynamics and population behavior*, Am. Nat., 149 (1997), pp. 247–269.
- [17] J. KENNEDY, S. KOÇAK, AND J. A. YORKE, *A chaos lemma*, Amer. Math. Monthly, 108 (2001), pp. 411–423.
- [18] U. KIRCHGRABER AND D. STOFFER, *On the definition of chaos*, Z. Angew. Math. Mech. 69 (1989), pp. 175–185.
- [19] S. A. LEVIN AND C. P. GOODYEAR, *Analysis of an age-structured fishery model*, J. Math. Biol., 9 (1980), pp. 245–274.
- [20] E. LIZ, *Global stability and bifurcations in a delayed discrete population model*, Int. J. Qual. Theory Differ. Equ. Appl., 3 (2009), pp. 66–80.
- [21] E. LIZ, *Complex dynamics of survival and extinction in simple population models with harvesting*, Theor. Ecol., 3 (2010), pp. 209–221.
- [22] E. LIZ AND D. FRANCO, *Global stabilization of fixed points using predictive control*, Chaos, 20 (2010), 023124, 9 pp.
- [23] E. LIZ AND A. RUIZ-HERRERA, *The hydra effect, bubbles, and chaos in a simple discrete population model with constant effort harvesting*, J. Math. Biol. (2012), available online, doi: 10.1007/s00285-011-0489-2

- [24] R.M. MAY, *Biological populations with nonoverlapping generations: stable points, stable cycles, and chaos*, Science, 186 (1974), pp. 645–647.
- [25] R.M. MAY, *Simple mathematical models with very complicated dynamics*, Nature, 261 (1976), pp. 459–467.
- [26] R.M. MAY, *Mathematical models in whaling and fisheries management*, in: G.F. Oster (ed.), *Some mathematical questions in biology*, Amer. Math. Soc., Providence, R.I., 1980, pp. 1–64.
- [27] A. MEDIO, M. PIREDDU, AND F. ZANOLIN, *Chaotic dynamics for maps in one and two dimensions: a geometrical method and applications to economics*, Internat. J. Bifur. Chaos Appl. Sci. Engrg., 19 (2009), pp. 3283–3309.
- [28] S. SMALE, *Finding a horseshoe on the beaches of Rio*, Math. Intelligencer 20 (1998), pp. 39–44.
- [29] H. R. THIEME, *Mathematics in population biology*, Princeton University Press, Princeton, 2003.
- [30] H. THUNBERG, *Periodicity versus chaos in one-dimensional dynamics*, SIAM Rev., 43 (2001), pp. 3–30.
- [31] S. TULJAPURKAR AND H. CASWELL (EDS.), *Structured-Population Models in Marine, Terrestrial, and Freshwater Systems*, Population and Community Biology Series, 18, Chapman & Hall, New York, 1997.
- [32] I. UGARCOVICI AND H. WEISS, *Chaotic dynamics of a nonlinear density dependent population model*, Nonlinearity, 17 (2004), pp. 1689–1711.
- [33] I. UGARCOVICI AND H. WEISS, *Chaotic attractors and physical measures for some density dependent Leslie population models*, Nonlinearity 20 (2007), pp. 2897–2906.
- [34] Q. WANG AND L.-S. YOUNG, *Strange attractors with one direction of instability*, Commun. Math. Phys 218 (2001), pp. 1–97.
- [35] A. WIKAN AND E. MJØLHUS, *Overcompensatory recruitment and generation delay in discrete age-structured population models.*, J. Math. Biol., 35 (1996), pp. 195–239.
- [36] A. A. YAKUBU, N. LI, J. M. CONRAD, AND M. L. ZEEMAN, *Constant proportion harvest policies: dynamic implications in the Pacific halibut and Atlantic cod fisheries*, Math. Biosci., 232 (2011), pp. 66–77.
- [37] P. ZGLICZYŃSKI AND M. GIDEA, *Covering relations for multidimensional dynamical systems*, J. Differential Equations, 202 (2004), pp. 32–58.
- [38] P. ZGLICZYŃSKI, *On periodic points for systems of weakly coupled 1-dim maps*. Nonlinear Anal., 46 (2001), pp. 1039–1062.