



Bootstrapping the statistical uncertainties of NN scattering data



R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola *

Departamento de Física Atómica, Molecular y Nuclear and Instituto Carlos I de Física Teórica y Computacional, Universidad de Granada, E-18071 Granada, Spain

ARTICLE INFO

Article history:

Received 16 July 2014

Received in revised form 18 September 2014

Accepted 18 September 2014

Available online 24 September 2014

Editor: W. Haxton

Keywords:

Monte Carlo simulation

NN interaction

One pion exchange

Statistical analysis

ABSTRACT

We use the Monte Carlo bootstrap as a method to simulate pp and np scattering data below pion production threshold from an initial set of over 6700 experimental mutually 3σ consistent data. We compare the results of the bootstrap, with 1020 statistically generated samples of the full database, with the standard covariance matrix method of error propagation. No significant differences in scattering observables and phase shifts are found. This suggests alternative strategies for propagating errors of nuclear forces in nuclear structure calculations.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

The modern era of high quality NN interactions started when the long term studies of the Nijmegen group culminated in a successful least squares fit with a statistically significant $\chi^2/\nu \sim 1$ [1] after implementation of many small but crucial effects and 3σ inconsistent data were excluded. Since then, subsequent analyses have been carried out [2–10] having $\chi^2/\nu \sim 1$ and with the purpose of being used in *ab initio* Nuclear Structure calculations. As is well known [11] any least-squares fit corresponds to χ^2 minimization

$$\min_{\mathbf{p}} \chi^2(\mathbf{p}) = \min_{\mathbf{p}} \sum_{i=1}^N \left(\frac{O_i^{\text{exp}} - O_i(\mathbf{p})}{\Delta O_i^{\text{exp}}} \right)^2 \equiv \chi^2(\mathbf{p}_0), \quad (1)$$

where O_i^{exp} is a fitted observable, ΔO_i^{exp} the corresponding statistical error bar and $O_i(\mathbf{p})$ the theoretical model depending on fitting parameters $\mathbf{p} = p_1, \dots, p_p$. The procedure assumes that the statistical uncertainties of the fitted data can be modeled by a probability distribution; namely independent normally distributed data $N(O_i^{\text{exp}}, \Delta O_i^{\text{exp}})$ an assumption based on counting a large number of events in NN scattering experiments. The assumption of a finite number of normally distributed data is an indispensable prerequisite for both meaningful uncertainty estimates from a

phenomenological fit and any subsequent and reliable error propagation. Fortunately, normality can be checked *a posteriori* in probabilistic terms and within a given confidence level by application of a variety of statistical tests, which naturally become more stringent with the number of data. Of course, individually checking the probability distribution of over 6700 data points, involving over 300 experiments, some dating back more than 60 years, is rather impractical. However, if a model fitted to the data is flexible enough to accurately reproduce them, the normality of the experimental data implies that discrepancies between theory and experiment, known as *residuals*, must follow a standard normal distribution, i.e.

$$R_i = \frac{O_i^{\text{exp}} - O_i(\mathbf{p}_0)}{\Delta O_i^{\text{exp}}} \sim N(0, 1). \quad (2)$$

Once a fit has been made, testing Eq. (2) is straightforward. Despite its simplicity, normality testing has not been a common practice in nuclear interactions fitting (an early discussion on normality was however conducted in Refs. [12,13]).

In a recent publication [9] we presented a new phenomenological Nucleon–Nucleon (NN) potential that accurately describes 6713 scattering data from 1950 to 2013 upgrading much of the previous works and increasing the statistics. This was done with an eye put on the determination of the uncertainties in the fitted NN interaction itself and their consequences in Nuclear Physics, for which little is still known (see however [14,15,6]) and statistical methods offer the most natural framework. On a more general level, a growing concern on the statistical analysis of nuclear theory and its predictive power has been initiated (see e.g. [16,17] for general

* Corresponding author.

E-mail addresses: rnavarrop@ugr.es (R. Navarro Pérez), amaro@ugr.es (J.E. Amaro), earriola@ugr.es (E. Ruiz Arriola).

and instructive overviews and references therein). We have applied some of the well known normality tests to three of our NN potentials, including the delta-shell potential with one pion exchange (DS-OPE) [8,9], (chiral) two pion exchange (DS- χ TPE) [10] and a gaussian potential with OPE [18] and found the normality condition to hold in all of them [18]. The lack of normality would clearly signal an inconsistency in the fitting analysis and might be used as a guide to unveil systematic errors both in the data as well as in the model. It is thus foreseeable that normality tests will be regarded as an important ingredient in the design of NN interactions statistically inferred from scattering data (see e.g. [19] for a posteriori analysis of [7]).

In our previous works the covariance matrix method was used to propagate errors. In the present note we discuss the robustness of our results using Monte Carlo techniques and the bootstrap method [20]. While these methods have successfully been exploited (see e.g. [21,22] for related studies within $\pi\pi$ scattering error analyses) to our knowledge they have never been implemented within the context of the NN force, so our presentation will be intentionally pedagogical. We also outline interesting consequences regarding strategies for error propagation in nuclear physics.

2. Covariance matrix method

In the standard covariance method one starts with a least squares fit Eq. (1). Once the condition of normality, Eq. (2), has been checked [18] and *assuming* normality of errors in the fitting parameters we are in position to propagate the statistical uncertainties into the potential parameters and any calculation that takes this potential parameters as an input. The error matrix \mathcal{E}_{ij} of the potential parameters $\{p_1, p_2, \dots, p_P\}$ can be calculated by inverting the Hessian matrix

$$H_{ij} = \left. \frac{\partial^2 \chi^2}{\partial p_i \partial p_j} \right|_{\mathbf{p}_0} \equiv (\mathcal{E}^{-1})_{ij} \quad (3)$$

which can be used to obtain confidence intervals for the parameters and correlations among them. Any quantity that can be calculated as a function of the potential parameters $F(p_1, p_2, \dots, p_P)$ can be provided with a statistical error bar ΔF with the customary expression

$$(\Delta F)^2 = \sum_{ij} \frac{\partial F}{\partial p_i} \frac{\partial F}{\partial p_j} \mathcal{E}_{ij}. \quad (4)$$

A good approximation to Eq. (3) can be found in [9] which has been used with Eq. (4) to estimate statistical uncertainties of phase-shifts, scattering amplitudes, deuteron properties, form factors, matrix elements and skyrme parameters [8–10,23,18,24]. However the derivatives in Eq. (4), depending on the functional form of F , may be hard to calculate analytically. If one contemplates numerical evaluation this requires a repeated evaluation of the function F at several values of the fitting parameters, which for a large number of parameters (typically 30–40 [8–10]) may also be a costly procedure.¹

¹ It can also be an inaccurate procedure since the corresponding finite differences step h must be smaller than the statistical Δp which are usually quite small. For instance, the evaluation of the Hessian numerically for our fits [8–10] requires to compute crossed derivatives, which turned out to be highly unstable for large number of parameters. This is why we preferred to compute the derivatives analytically and use in passing the highly efficient Levenberg–Marquardt minimization algorithm where a stable (definite positive) approximation to the Hessian is exploited [25].

The calculation of derivatives can be avoided by drawing random numbers following a multivariate normal distribution determined by the covariance matrix \mathcal{E} ,

$$P(p_1, p_2, \dots, p_P) = \frac{1}{\sqrt{(2\pi)^P \det \mathcal{E}}} e^{-\frac{1}{2}(\mathbf{p}-\mathbf{p}_0)^T \mathcal{E}^{-1}(\mathbf{p}-\mathbf{p}_0)}. \quad (5)$$

This generates a family of potential parameters and calculate F with each potential. This Monte Carlo method directly propagates uncertainties, however a multivariate normal probability distribution to all the parameters is assumed which may not always be the case for the *true* distribution of parameters.

3. The bootstrap method

The Bootstrap is a Monte Carlo technique that allows to find the most likely parameters probability distribution and propagate statistical uncertainties and correlations into any function F [20] (see also [25]). In our case the deviations between the theoretical model and the experimental data are normal statistical fluctuations. The procedure corresponds to generate replicas of the observed data which are meant to simulate a fictitious experiment. Thus, for every experimental data point O_i^{exp} with uncertainty ΔO_i^{exp} one generates M “synthetic” random points $O_{i,1}^{\text{synth}}, O_{i,2}^{\text{synth}}, \dots, O_{i,M}^{\text{synth}}$ distributed as $N(O_i^{\text{exp}}, \Delta O_i^{\text{exp}})$, i.e.

$$O_{i,\alpha}^{\text{synth}} = O_i^{\text{exp}} + \xi_{i,\alpha} \Delta O_i^{\text{exp}} \quad (6)$$

where $\xi_{i,\alpha} \sim N(0, 1)$ are standard normal and independent variables, $\langle \xi_{i,\alpha} \rangle = 0$ and $\langle \xi_{i,\alpha} \xi_{j,\beta} \rangle = \delta_{ij} \delta_{\alpha\beta}$. This will generate M independent databases with the same number of data as the original one. Each synthetic database will represent a snapshot of the random fluctuations inherent to the experimental processes. A least squares fit to every generated database, $O_{i,\alpha}^{\text{synth}}$ ($\alpha = 1, \dots, M$), featuring a maximum likelihood estimate can be made and a family of parameters $p_{1,\alpha}, \dots, p_{P,\alpha}$ will be obtained as

$$\min_{\mathbf{p}} \chi_\alpha^2(\mathbf{p}) = \min_{\mathbf{p}} \sum_{i=1}^N \left(\frac{O_{i,\alpha}^{\text{synth}} - O_i(\mathbf{p})}{\Delta O_i^{\text{exp}}} \right)^2 \equiv \chi^2(\mathbf{p}_\alpha). \quad (7)$$

Then, the most likely theory parameters are \mathbf{p}_α . The corresponding joined or marginal probability distributions can be obtained by binning the outcoming parameter samples. This allows to compute any function of the theoretical model parameters $F(\mathbf{p})$ at a set of points $F_\alpha \equiv F(\mathbf{p}_\alpha)$. Thus, the mean and variance can be computed for large M as usual,

$$E(F) = \frac{1}{M} \sum_{\alpha=1}^M F(\mathbf{p}_\alpha), \quad (8)$$

$$\text{Var}(F) = \frac{M}{M-1} E[(F - E(F))^2], \quad (9)$$

The correlation coefficient of two different observables is

$$\mathcal{C}(F, G) = \frac{E[(F - E(F))(G - E(G))]}{\sqrt{E[(F - E(F))^2]} \sqrt{E[(G - E(G))^2]}}, \quad (10)$$

so that $\mathcal{C}(p_i, p_j) = \mathcal{C}_{ij} = \mathcal{E}_{ij}/(\mathcal{E}_{ii}\mathcal{E}_{jj})^{\frac{1}{2}}$ is the correlation matrix. For asymmetric or skewed distributions, it may be better to define the 1σ asymmetric coverage by excluding 16% of the upper and lower values of the distribution instead of the variance definition, Eq. (9). At any rate we always check this possibility before errors are quoted.

While the bootstrap method requires to perform M repeated fits, it is a competitive alternative to determine errors and correlations when the covariance matrix itself is not directly available nor




Fig. 1. (Color online.) Correlation matrix \mathcal{C}_{ij} for the DS-OPE potential parameters $(\lambda_{i,l})_{l,l'}^S$ in the partial wave basis [8]. The points $r_i = \Delta r(i+1)$ are grouped within every partial wave. We show the results obtained with the covariance matrix (left panel) and the Monte Carlo bootstrap simulation of experimental data (right panel). We grade gradually from 100% correlation, $\mathcal{C}_{ij} = 1$ (red), 0% correlation, $\mathcal{C}_{ij} = 0$ (yellow) and 100% anti-correlation, $\mathcal{C}_{ij} = -1$ (blue).

used in the minimization method [26]. Again, we stress that this method to generate snapshots of the statistical fluctuations is justified since the condition of Eq. (2) has been checked to a significant confidence level.

4. Numerical results

We apply the different methods to the 3σ self-consistent database presented in Ref. [9] where $\chi^2/\nu = 1.04$. The potential used for this analysis has the form

$$V(\vec{r}) = V_{\text{short}}(r)\theta(r_c - r) + V_{\text{long}}(r)\theta(r - r_c). \quad (11)$$

The long range piece $V_{\text{long}}(\vec{r})$ contains a Charge-Dependent (CD) One pion exchange (OPE) with a fixed $f^2 = 0.075$ [13] and electromagnetic (EM) corrections which are kept fixed throughout the fitting process. The short component was inspired by Avilés [27] (see also [28,29]) and reads

$$V_{\text{short}}(\vec{r}) = \sum_{n=1}^{21} \hat{O}_n \left[\sum_{i=1}^N V_{i,n} \delta(r - r_i) \right], \quad (12)$$

where \hat{O}_n are the set of operators in the extended AV18 basis [3,14,15,23], $V_{i,n}$ are fitting parameters and $r_i = \Delta r(i+1)$ with $\Delta r = 0.6\text{fm}$. The fit is carried out more effectively in terms of some low and independent partial waves contributions to the potential $(\lambda_{i,\alpha})_{l,l'}^S$ from which all other higher partial waves are consistently deduced (see Ref. [8,9]). The delta-shell potential reduces the computational effort enormously, so a large number of fits can easily be undertaken.

For the bootstrap analysis we took $M = 1020$ samples of the $N = 6713$ data and refitted the parameters of the DS-OPE potential (denoted by $(\lambda_{i,\alpha})_{l,l'}^S$) which was used to determine the database. This generates M independent sets of most likely parameters to each synthetic database $O_{i,\alpha}$, $\alpha = 1, \dots, M$. From there any function of the fitted parameters and the inherent correlations can be determined.

In Fig. 1 we show the correlation matrix of the DS-OPE potential parameters obtained with the standard covariance matrix method and the Bootstrap method. It is not obvious, though most welcome, that both covariance and bootstrap methods give fairly similar results, although small correlations are overestimated by the covariance matrix. The main difference between both methods is in the 3S_1 - 3D_1 coupled channel. The Monte Carlo bootstrap

simulation results in very small correlations between the 3S_1 and ϵ_1 partial wave parameters and stronger correlations between ϵ_1 and 3D_1 ; in contrast the covariance matrix method gives opposite results. These discrepancies could be related to the fitting of the deuteron binding energy where the approximation used for the Hessian matrix might be outside of its range of validity. In fact, the Monte Carlo generated 3S_1 , ϵ_1 and 3D_1 parameters show large asymmetries as can clearly be seen in Fig. 2.

We compare in Fig. 3 the propagation of statistical uncertainties into phase-shifts by three methods: i) the standard covariance matrix method, ii) the equivalent Monte Carlo implementation of the covariance matrix using the multivariate normal distribution of Eq. (5) with $M = 1020$, and iii) the bootstrap method also with $M = 1020$ samples. The first and the second methods should produce the same results for a sufficiently large number of parameter samples. So the agreement between Eq. (4) and the Monte Carlo sampling of parameters Eq. (5) reflects the large M value with the same \mathcal{E} . Although the bootstrap method tends to give slightly larger error bars the difference with the other two methods is not significant. As mentioned above, one potential advantage of the Bootstrap method is that it relaxes the assumption of normally distributed fitted parameters, a feature which proves relevant for asymmetric or skewed distributions. We find that the asymmetries seen in Fig. 2 do not significantly propagate to the corresponding phase shifts.

As a matter of principle the Monte Carlo simulation of data gives the most reliable uncertainty propagation, but considering that performing a large number of full-length fits to data can be computationally expensive the covariance matrix methods are a fairly good and extremely useful approximation which will be exploited in future work.

5. Conclusions

The propagation of statistical errors of nuclear forces stemming from the finite precision and number of experimental NN scattering data requires in the first place passing a normality test. However, even in this favourable case the actual calculation may be computationally demanding because of a practical need of repeating large scale computations. It is thus important to explore methods where the number of calculations can be kept to a minimum. In the standard covariance matrix method one needs the evaluation of the Hessian as well as the derivatives of the object function whose uncertainties




Fig. 2. (Color online.) 3S_1 - 3D_1 coupled channel Delta Shell parameters distribution. The parameters are 3S_1 partial wave (upper row), ϵ_1 mixing angle (middle row) and 3D_1 partial wave (lower row). The blue bars give the normalized histogram from the 1020 fits to the Monte Carlo generated databases. The red line is the normal distribution given to each parameter by the covariance matrix method.




Fig. 3. (Color online.) Low angular momentum partial wave phase-shifts statistical error bars calculated from the DS-OPE potential using the covariance matrix with Eq. (4) (left panel), a Monte Carlo sample of the potential parameters according to the same covariance matrix (middle panel) and the collection of parameters resulting from bootstrapping the database (right panel).

are evaluated with respect to the theoretical model parameters. As an alternative the Monte Carlo method based on explicit knowledge of the Hessian can profitably be used as it avoids the computation of derivatives (analytical or numerical) and automatically implements in any snapshot the inherent correlations in the fitting parameters. The previous methods assume a multivariate normal distribution of the fitting parameters.

We have thus analyzed the more elaborated bootstrap method which also rests on the normality test and is based on a multiple minimization to a synthetic set of data generated by the distribution of the most likely estimate of the model parameters. While this approach assumes normality of the experimental data but not of the fitting parameters, it allows to handle possible skewness in the parameter distributions. Our bootstrap analysis confirms the error and correlations already found by the covariance method.

Acknowledgements

This work is supported by Spanish DGI (grant FIS2011-24149) and Junta de Andalucía (grant FQM225). R.N.P. is supported by a Mexican CONACYT fellowship.

References

- [1] V.G.J. Stoks, R.A.M. Klomp, M.C.M. Rentmeester, J.J. de Swart, *Phys. Rev. C* 48 (1993) 792.
- [2] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen, J.J. de Swart, *Phys. Rev. C* 49 (1994) 2950, arXiv:nucl-th/9406039.
- [3] R.B. Wiringa, V. Stoks, R. Schiavilla, *Phys. Rev. C* 51 (1995) 38, arXiv:nucl-th/9408016.
- [4] R. Machleidt, *Phys. Rev. C* 63 (2001) 024001, arXiv:nucl-th/0006014.
- [5] F. Gross, A. Stadler, *Phys. Rev. C* 78 (2008) 014005, arXiv:0802.1552.
- [6] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Phys. Lett. B* 724 (2013) 138, arXiv:1202.2689.
- [7] A. Ekström, G. Baardsen, C. Forssén, G. Hagen, M. Hjorth-Jensen, et al., *Phys. Rev. Lett.* 110 (2013) 192502, arXiv:1303.4674.
- [8] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Phys. Rev. C* 88 (2013) 024002, arXiv:1304.0895.
- [9] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Phys. Rev. C* 88 (2013) 064002, arXiv:1310.2536.
- [10] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Phys. Rev. C* 89 (2014) 024004, arXiv:1310.6972.
- [11] M.J. Evans, J.S. Rosenthal, *Probability and Statistics: The Science of Uncertainty*, Macmillan, 2004.
- [12] J. Bergervoet, P. van Campen, W. van der Sanden, J.J. de Swart, *Phys. Rev. C* 38 (1988) 15.
- [13] V.G. Stoks, R. Timmermans, J. de Swart, *Phys. Rev. C* 47 (512) (1993), arXiv:nucl-th/9211007.
- [14] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, arXiv:1202.6624, 2012.
- [15] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *PoS QNP2012* (2012) 145, arXiv:1206.3508, 2012.
- [16] J. Dudek, B. Szpak, B. Fornal, A. Dromard, *Phys. Scr.* 2013 (2013) 014002.
- [17] J. Dobaczewski, W. Nazarewicz, P.G. Reinhard, *J. Phys. G, Nucl. Part. Phys.* 41 (2014) 074001.
- [18] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Phys. Rev. C* 89 (2014) 064006, arXiv:1404.0314.
- [19] A. Ekström, B. Carlsson, K. Wendt Forssén, M. Hjorth-Jensen, et al., arXiv:1406.6895, 2014.
- [20] B. Efron, *The Jackknife, the Bootstrap and Other Resampling Plans*, vol. 38, SIAM, 1982.
- [21] J. Nieves, E. Ruiz Arriola, *Eur. Phys. J. A* 8 (2000) 377, arXiv:hep-ph/9906437.
- [22] J. Nieves, E. Ruiz Arriola, *Nucl. Phys. A* 679 (2000) 57, arXiv:hep-ph/9907469.
- [23] J.E. Amaro, R. Navarro Pérez, E. Ruiz Arriola, *Few-Body Syst.* 55 (2014) 977.
- [24] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *J. Phys. G, Nucl. Part. Phys.*, in press, arXiv:1406.0625, 2014.
- [25] W. Press, S. Teukolsky, W. Vetterling, B. Flannery, *Numerical Recipes in FORTRAN 2*, vol. 77, Cambridge University Press, New York, 2007.
- [26] M. Kortelainen, T. Lesinski, J. More, W. Nazarewicz, J. Sarich, et al., *Phys. Rev. C* 82 (2010) 024313, arXiv:1005.5145.
- [27] J. Aviles, *Phys. Rev. C* 6 (1972) 1467.
- [28] D. Entem, E. Ruiz Arriola, M. Pavon Valderrama, R. Machleidt, *Phys. Rev. C* 77 (2008) 044006, arXiv:0709.2770.
- [29] R. Navarro Pérez, J.E. Amaro, E. Ruiz Arriola, *Prog. Part. Nucl. Phys.* 67 (2012) 359, arXiv:1111.4328.