

THE RELATIVE ISOPERIMETRIC INEQUALITY OUTSIDE CONVEX DOMAINS IN \mathbf{R}^n

JAIGYOUNG CHOE, MOHAMMAD GHOMI, AND MANUEL RITORÉ

ABSTRACT. We prove that the area of a hypersurface Σ which traps a given volume outside a convex domain C in Euclidean space \mathbf{R}^n is bigger than or equal to the area of a hemisphere which traps the same volume on one side of a hyperplane. Further, when C has smooth boundary ∂C , we show that equality holds if and only if Σ is a hemisphere which meets ∂C orthogonally.

1. INTRODUCTION

Let $\mathbf{H}^n := \{(x_1, \dots, x_n) \in \mathbf{R}^n : x_n \geq 0\}$ be the closed upper half of Euclidean space \mathbf{R}^n . Given $D \subset \mathbf{H}^n$, reflection across the boundary $\partial\mathbf{H}^n$ and the classical isoperimetric inequality in \mathbf{R}^n imply that

$$\left(\text{area}(\partial D \sim \partial\mathbf{H}^n)\right)^n \geq \frac{1}{2} n^n \omega_n (\text{vol } D)^{n-1},$$

with equality if and only if D is a half ball and $\partial D \sim \partial\mathbf{H}^n$ is a hemisphere. Here *area* and *vol* (*volume*) denote, respectively, the $(n-1)$ and n dimensional Hausdorff measures, ω_n is the volume of the unit ball in \mathbf{R}^n , and \sim is the set exclusion operator. In this paper we prove that the above inequality holds outside any convex set $C \subset \mathbf{R}^n$ with interior points, i.e.,

$$(1.1) \quad \left(\text{area}(\partial D \sim \partial C)\right)^n \geq \frac{1}{2} n^n \omega_n (\text{vol } D)^{n-1},$$

for any $D \subset \mathbf{R}^n \sim C$. Further we show that when ∂C is smooth (C^∞), equality holds if and only if D is a half ball and $\partial D \sim \partial C$ is a hemisphere.

We call (1.1) the *relative isoperimetric inequality of D* with supporting set C . The proof of this inequality for $n = 2$ is easy once one reflects the convex hull of D about its linear boundary. For $n \geq 3$ some partial results were known: I. Kim [11] proved (1.1) for $C = U \times \mathbf{R}$, where U is the epigraph of a C^2

Date: March 13, 2006.

2000 *Mathematics Subject Classification.* 58E35, 49Q20.

Work of M. G. was supported in part by NSF Grant DMS-0336455, and CAREER award DMS-0332333. Work of J. C. was supported in part by KOSEF R01-2000-000-00007-0. Work of J. C. and M. R. was supported in part by MEC research project MTM2004-01387.

convex function, and J. Choe [2] proved (1.1) when $\partial D \cap \partial C$ is a graph which is symmetric about $(n - 1)$ hyperplanes of \mathbf{R}^n . More recently, J. Choe and M. Ritoré [4] have shown that (1.1) holds outside convex sets in 3-dimensional Cartan-Hadamard manifolds, with equality if and only if D is a flat half ball and $\Sigma := \partial D \sim \partial C$ is a hemisphere. The main ingredients of the proof in [4] are the estimate $(\sup_{\Sigma} H^2) \text{ area } \Sigma \geq 2\pi$, and the analysis of the equality case, where H is the mean curvature of Σ ; however, the methods used in [4], which were inspired by the work of P. Li and S.-T. Yau [12], are valid only when $n = 3$.

We obtain inequality (1.1), and the characterization of its equality case presented below, from the estimate

$$\left(\sup_{\Sigma} H^{n-1} \right) \text{ area } \Sigma \geq \frac{\mathbf{c}_{n-1}}{2},$$

where \mathbf{c}_{n-1} is the area of the unit sphere $\mathbf{S}^{n-1} \subset \mathbf{R}^n$. This inequality follows from the arithmetic-geometric mean inequality between H (the average of the principal curvatures) and the Gauss-Kronecker curvature GK (the product of principal curvatures) of Σ , once we show that the total Gauss-Kronecker curvature of the set of regular points of $\partial D \sim \partial C$ with positive principal curvatures is larger than or equal to $\mathbf{c}_{n-1}/2$. We proved the latter inequality in [3] assuming slightly more regularity than is warranted in the present case; however, as we verify below, that proof essentially works here as well.

2. PRELIMINARIES: EXISTENCE AND REGULARITY

Throughout this paper $C \subset \mathbf{R}^n$ denotes a *proper convex set*, which we define as a closed convex set with interior points and nonempty boundary ∂C . Further, unless noted otherwise we assume that ∂C is \mathcal{C}^∞ , which is what we mean when we say that C has *smooth* boundary. For any $A \subset \mathbf{R}^n$, let $A_C := A \sim C$. The *relative isoperimetric profile* of \mathbf{R}_C^n is the function $I_C : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ given by

$$I_C(v) := \inf_D \{ \text{area}(\partial D)_C : D \Subset \mathbf{R}^n, \text{ vol } D = v \},$$

where $D \Subset \mathbf{R}^n$ means that D is relatively compact in \mathbf{R}^n . Note that

$$I_{\mathbf{H}^n}(v) = n \left(\frac{\omega_n}{2} \right)^{\frac{1}{n}} v^{\frac{n-1}{n}}.$$

So the relative isoperimetric inequality (1.1) is equivalent to

$$(2.1) \quad I_C(\text{vol } D) \geq I_{\mathbf{H}^n}(\text{vol } D).$$

An *isoperimetric region* $D \subset \mathbf{R}_C^n$ is one for which the equality $\text{area}(\partial D)_C = I_C(\text{vol } D)$ holds. An isoperimetric region need not exist for a given volume.

Denote by $\mathcal{C}_0^1(\mathbf{R}_C^n, \mathbf{R}^n)$ the set of \mathcal{C}^1 vector fields with compact support in \mathbf{R}_C^n . For any $D \subset \mathbf{R}^n$, the *perimeter* of D relative to C is defined as

$$\mathcal{P}_C(D) := \sup \left\{ \int_D \operatorname{div} X : X \in \mathcal{C}_0^1(\mathbf{R}_C^n, \mathbf{R}^n), |X| \leq 1 \right\},$$

where $|X|$ is the supremum norm. Stokes theorem implies that the perimeter of a set and the area of its boundary coincide for sets whose boundary is a \mathcal{C}^1 hypersurface (see Giusti's book [6] for background on finite perimeter sets). In order to prove (1.1), we need to minimize \mathcal{P}_C subject to a volume constraint, i.e., given $v \in (0, \operatorname{vol} E)$, we wish to find $\Omega_0 \subset E$, with $\operatorname{vol} \Omega_0 = v$, such that

$$\mathcal{P}_C(\Omega_0) \leq \mathcal{P}_C(\Omega),$$

for any $\Omega \subset E$ with $\operatorname{vol} \Omega = v$. The existence of Ω_0 is guaranteed by the boundedness of E , see [6], and the regularity properties of Ω_0 which we need may be summarized as follows

Lemma 2.1. *Let E be the closure of a bounded domain with smooth boundary in \mathbf{R}_C^n . Then, for any $v \in (0, \operatorname{vol} E)$, there is a set $\Omega_0 \subset E$ of volume v minimizing \mathcal{P}_C . Moreover*

- (i) ([7]) $\partial\Omega_0$ has constant mean curvature and is smooth in the interior of E except for a singular set of Hausdorff dimension less than or equal to $(n - 8)$.
- (ii) ([9, p. 263]) $\overline{(\partial\Omega_0)_C}$ meets ∂C orthogonally except for a singular set of Hausdorff dimension less than or equal to $(n - 8)$. In fact $\overline{(\partial\Omega_0)_C}$ is smooth at every point of $\overline{(\partial\Omega_0)_C} \cap \partial C$ away from this singular set.
- (iii) ([15, Thm. 3.6]) If $(\partial E)_C$ is strictly convex then $(\partial\Omega_0)_C$ meets $(\partial E)_C$ tangentially and it is $\mathcal{C}^{1,1}$ in a neighborhood of $(\partial E)_C$.
- (iv) At every point $x_0 \in (\partial\Omega_0)_C$ there is a tangent cone obtained by blowing up the set Ω_0 about x_0 . If this tangent cone is contained in a half space of \mathbf{R}^n , then it is the half space and $(\partial\Omega_0)_C$ is regular at x_0 [6]. At points in $\overline{(\partial\Omega_0)_C} \cap \partial C \sim \overline{(\partial E)_C}$ we have the same result, as described in [9].

The $\mathcal{C}^{1,1}$ regularity of $(\partial\Omega_0)_C$ near $(\partial E)_C$ will be enough for our purposes here since, by Rademacher's Theorem, a $\mathcal{C}^{1,1}$ hypersurface has principal curvatures defined almost everywhere, and thus we will be able to apply the integral curvature estimates obtained in the next two sections.

3. THE ESTIMATE FOR TOTAL POSITIVE CURVATURE

First we recall the general definition for total positive curvature τ^+ studied in [3]. Let $\Sigma = (\Sigma \sim \Sigma_0) \cup \Sigma_0$ be the compact union of a $\mathcal{C}^{1,1}$ hypersurface

$\Sigma \sim \Sigma_0$ with boundary and a singular set Σ_0 of Hausdorff dimension less than or equal to $(n - 8)$ so that $\Sigma_0 \subset \overline{\Sigma \sim \Sigma_0}$. The points in $\Sigma \sim \Sigma_0$ will be called *regular points* of Σ . A hyperplane $\Pi \subset \mathbf{R}^n$ is called a *restricted support hyperplane* of Σ at a point p , if $p \in \Pi \cap \Sigma$, Σ lies on one side of Π , and Π is tangent to Σ when $p \in \partial\Sigma \sim \Sigma_0$. An *outward normal* of Π is a normal vector to Π which points towards a side of Π not containing Σ . If Π is a restricted support hyperplane for an open neighborhood U_p of p in Σ , then Π is called a *restricted local support hyperplane*; furthermore, p is a *locally strictly convex point* of Σ , or $p \in \Sigma^+$, provided that $\Pi \cap U_p = \{p\}$. Let $N(\Sigma^+) \subset \mathbf{S}^{n-1}$ be the set of outward unit normals to restricted local support hyperplanes of Σ at points of Σ^+ . Then the *total positive curvature* τ^+ of Σ is defined as the algebraic area (i.e. area counted with multiplicity) of $N(\Sigma^+)$, where by *area* we mean the $(n - 1)$ -dimensional Hausdorff measure \mathcal{H}^{n-1} . More formally, if for each $u \in N(\Sigma^+)$, we let $\Sigma_u^+ \subset \Sigma^+$ be the set of points where Σ has a restricted local support hyperplane with outward unit normal u , then

$$\tau^+(\Sigma) := \int_{u \in N(\Sigma^+)} d\mathcal{H}^{n-1}(\Sigma_u^+)$$

where integration is with respect to the volume element or the $(n-1)$ -dimensional Hausdorff measure on \mathbf{S}^{n-1} .

As we are assuming that $\Sigma \sim \Sigma_0$ is a $\mathcal{C}^{1,1}$ hypersurface, the principal curvatures are defined for almost every point of $\Sigma \sim \Sigma_0$ and so the Gauss-Kronecker curvature GK , the product of all principal curvatures, may be integrated on $\Sigma \sim \Sigma_0$. Moreover, in case there are no restricted local support hyperplanes of Σ at points of Σ_0 , the *area formula* [5, Thm 3.2.3] yields that

$$\tau^+(\Sigma) = \int_{\Sigma^+ \sim \Sigma_0} GK.$$

As remarked in the introduction, the main ingredient in the proof of the relative isoperimetric inequality is the following estimate. We state below the version for convex sets with smooth boundary we shall need. The proof of this result is a slight modification of the one given in the appendix of [3].

Lemma 3.1. *Let $\Sigma = (\Sigma \sim \Sigma_0) \cup \Sigma_0 \subset \mathbf{R}^n$ be the union of a $\mathcal{C}^{1,1}$ embedded hypersurface $\Sigma \sim \Sigma_0$ and a singular set Σ_0 such that $\partial\Sigma \sim \Sigma_0$ is a \mathcal{C}^2 submanifold that lies on the boundary of a convex set $C \subset \mathbf{R}^n$ with \mathcal{C}^2 boundary ∂C . Suppose that there are no restricted local support hyperplanes of Σ at points of Σ_0 , and that, at each point $p \in \partial\Sigma \sim \Sigma_0$, the inward conormal $\sigma(p)$ of $\partial\Sigma$ is the outward unit normal to C at p . Then*

$$(3.1) \quad \tau^+(\Sigma) \geq \frac{\mathbf{c}_{n-1}}{2},$$

and equality holds if and only if $\partial\Sigma$ lies in a hyperplane.

Proof. Let $\partial\Sigma_r := \partial\Sigma \sim \Sigma_0$ be the regular part of the boundary of Σ ,

$$U\partial\Sigma_r := \{ (p, u) \mid p \in \partial\Sigma_r, u \in \mathbf{S}^{n-1}, u \perp T_p\partial\Sigma_r \}$$

be the *unit normal bundle* of $\partial\Sigma_r$, and $\nu: U\partial\Sigma_r \rightarrow \mathbf{S}^{n-1}$, given by

$$\nu(p, u) := u,$$

be its *Gauss map*. Define $I_r \subset J_r \subset U\partial\Sigma_r$ by

$$\begin{aligned} I_r &:= \{ (p, u) \in U\partial\Sigma_r \mid \langle x - p, u \rangle \leq 0, \quad \forall x \in \Sigma \}, \\ J_r &:= \{ (p, u) \in U\partial\Sigma_r \mid \langle x - p, u \rangle \leq 0, \quad \forall x \in \partial\Sigma \}. \end{aligned}$$

Note that if $(p, u) \in J_r \sim I_r$, then the height function $x \mapsto \langle x - p, u \rangle$ achieves its maximum in the interior of Σ , and thus Σ has a restricted support hyperplane with outward normal u . By hypothesis, the point p must then lie in the regular part $\Sigma \sim \Sigma_0$ of Σ . Hence

$$\tau^+(\Sigma) \geq \text{area } \nu(J_r \sim I_r),$$

since almost every support hyperplane of Σ intersects Σ at a single point [14, Thm. 2.2.9]. So to prove (3.1) it suffices to show that

$$\text{area } \nu(J_r \sim I_r) \geq \frac{\mathbf{c}_{n-1}}{2}.$$

The proof of the last inequality is virtually identical to the proof of the corresponding inequality (9) in the appendix of [3] to which we refer the reader. \square

4. THE MEAN CURVATURE ESTIMATE

As we will see in the next section, in order to prove (1.1), we need to construct a bounded region E outside C in \mathbf{R}^n , and minimize the perimeter \mathcal{P}_C under a volume constraint inside E . We shall see in our next result that the boundary of any isoperimetric region so obtained satisfies the hypotheses of Lemma 3.1. In particular, the lower curvature bound (3.1) holds for such regions, which in turn yields the following estimate for mean curvature.

Proposition 4.1. *Let $C \subset \mathbf{R}^n$ be a proper convex set, $p_0 \in \partial C$, $E := \overline{B(p_0, r)}_C$, $\Omega \subset E_C$ be a set minimizing the perimeter \mathcal{P}_C under a volume constraint, and H_Σ be the (constant) mean curvature of the regular part of $\Sigma := \overline{(\partial\Omega)_C}$. Then*

$$(4.1) \quad H_\Sigma^{n-1} \mathcal{P}_C(\Omega) \geq \frac{\mathbf{c}_{n-1}}{2}.$$

Equivalently, if $H_0(a)$ denotes the mean curvature of a hemisphere of area a , then

$$H_\Sigma \geq H_0(\mathcal{P}_C(\Omega)).$$

Equality holds in these inequalities if and only if Ω is a half ball and Σ meets ∂C orthogonally.

Proof. It is enough to show that, if Π is a support hyperplane of Ω at $p \in \Sigma$, then p is a regular point of Σ .

If $p \in \Sigma \cap \text{int}(E)$ then the minimal tangent cone of Ω at p is contained in a half space. By [6, Thm. 15.5], it must be a half space and so Σ is regular at p .

If $p \in \partial\Sigma \sim \overline{(\partial E)_C}$ then we consider the integer multiplicity rectifiable current $\partial[\Omega]$. Reflecting it [8, Remark 3.1] with respect to ∂C and blowing up about p we get an area-minimizing oriented tangent cone T [8, Thm. 3.5], [9]. Let H be the tangent hyperplane of ∂C at p , and H^+ the closed half space determined by H whose interior does not meet C . Assuming there is a support hyperplane Π of Ω at p , we get that the support of T , $\text{supp}(T)$, is contained in a region of \mathbf{R}^n bounded by $H_1 \cup H_2$, where $H_1 = \Pi \cap H^+$ and H_2 is the reflection of H_1 with respect to H . Let $S = H_1 \cap H = H_1 \cap H_2$. We have that S is an $(n-2)$ -dimensional linear submanifold of \mathbf{R}^n which is contained in H .

Rotating H_1, H_2 with respect to S until they first touch $\text{supp}(T) \sim S$, using the maximum principle, and a connectedness argument, we get that $\text{supp}(T) = H_1 \cup H_2$, which is not area-minimizing unless $H_1 \cup H_2$ is a hyperplane orthogonal to H . Hence Σ is regular at p .

Observe that $\partial\Sigma \cap \partial C \cap \overline{(\partial E)_C} = \emptyset$: if $x_0 \in \partial\Sigma \cap \partial C \cap \overline{(\partial E)_C}$, then the outer normal ν to ∂C and the outer normal $\tilde{\nu}$ to $\partial B(p, r)$ satisfy $\langle \nu, \tilde{\nu} \rangle(x_0) > 0$. Reasoning as in the two previous paragraphs, reflecting and blowing up about x_0 we get a cone which minimizes area in a wedge of angle less than π , thus getting a contradiction.

So we can apply Lemma 3.1 to conclude that

$$\int_{\Sigma^+ \sim \Sigma_0} GK = \tau^+(\Sigma) \geq \frac{c_{n-1}}{2}.$$

By [15, Thm. 3.7], H_Σ , the constant mean curvature of the regular part $\Sigma \sim \Sigma_0$ of Σ in the interior of E , is an upper bound for the mean curvature of Σ . So we have

$$H_\Sigma^{n-1} \mathcal{P}_C(\Omega) \geq \int_{\Sigma^+ \sim \Sigma_0} H_\Sigma^{n-1} \geq \int_{\Sigma^+ \sim \Sigma_0} GK \geq \frac{c_{n-1}}{2},$$

which establishes the first desired inequality. To obtain the second inequality note that, if r is the radius of a hemisphere of area $\mathcal{P}_C(\Omega)$, then

$$\left(H_0(\mathcal{P}_C(\Omega))\right)^{n-1} \mathcal{P}_C(\Omega) = \left(\frac{1}{r}\right)^{n-1} \frac{\mathbf{c}_{n-1} r^{n-1}}{2} = \frac{\mathbf{c}_{n-1}}{2} \leq H_{\Sigma}^{n-1} \mathcal{P}_C(\Omega).$$

If equality holds then $\Sigma \sim \Sigma_0 = \Sigma^+$, and $H_{\Sigma}^{n-1} = GK$, which implies that $\Sigma \sim \Sigma_0$ is totally umbilical and so Σ_0 is empty. Further $\partial\Sigma$ lies in a hyperplane by Lemma 3.1, and so Ω is a half ball and Σ intersects ∂C orthogonally. \square

5. PROOF OF THE RELATIVE ISOPERIMETRIC INEQUALITY IN \mathbf{R}_C^n

With the aid of Proposition 4.1 we are now in a position to prove the main result of this paper.

Theorem 5.1. *Let $C \subset \mathbf{R}^n$ be a proper convex set (with smooth boundary). For any bounded set $D \subset \mathbf{R}_C^n$ with finite perimeter,*

$$(5.1) \quad (\text{area}(\partial D)_C)^n \geq \frac{1}{2} n^n \omega_n (\text{vol } D)^{n-1},$$

with equality if and only if D is a half ball and $(\partial D)_C$ is a hemisphere.

Remark 5.2. If C is bounded then, from the results in [13], it can be proved that any perimeter minimizing sequence of sets in \mathbf{R}_C^n of given volume has a subsequence converging to an isoperimetric region. In this case the proof of Theorem 5.1 can be slightly simplified. However, when C is unbounded, we have to deal with the possibility of nonexistence of minimizers in \mathbf{R}_C^n .

Proof of Theorem 5.1. First we construct an exhaustion of \mathbf{R}_C^n . Fix $p_0 \in \partial C$, and let $\{r_m\}_{m \in \mathbb{N}}$ be a diverging sequence of positive increasing numbers. In case C is bounded we require that $C \subset B(p_0, r_m)$. We define $E_m := \overline{B(p_0, r_m)}_C$.

Since E_m is bounded, isoperimetric regions exist in E_m for any given volume $v \in (0, \text{vol } E_m)$. Let $\Omega \subset E_m$ be an isoperimetric region minimizing \mathcal{P}_C in E_m under a volume constraint, and let $\Sigma := \overline{(\partial\Omega)}_C$.

By Proposition 4.1, for every component Ω' of Ω touching the boundary of C , with $\Sigma' := \overline{(\partial\Omega')}_C$, we have

$$H_{\Sigma'}^{n-1} \mathcal{P}_C(\Sigma') \geq \frac{\mathbf{c}_{n-1}}{2},$$

with equality if and only if Ω' is an open half ball and Σ' is an open hemisphere. Observe that, for a component Ω'' of Ω not touching the boundary of C , with $\Sigma'' := \overline{(\partial\Omega'')}_C$, one easily obtains

$$H_{\Sigma''}^{n-1} \mathcal{P}_C(\Sigma'') \geq \mathbf{c}_{n-1},$$

with equality if and only if Ω'' is a ball and Σ'' a round sphere.

Breaking Ω into components touching ∂C and components in the interior of \mathbf{R}_C^n we get

$$H_\Sigma^{n-1} \mathcal{P}_C(\Omega) \geq \frac{c_{n-1}}{2},$$

and equality holds if and only if Ω consists of one connected component which is a half ball, and Σ an open hemisphere.

Let I_m be the isoperimetric profile of E_m . From standard arguments, see [10, p. 170–172,], it follows that (i) I_m is continuous and increasing, (ii) if I_m is smooth at v_0 , then $I'_m(v_0) = (n-1)H$, where H is the constant mean curvature in the interior of E_m of *any* isoperimetric region of volume v_0 , and (iii) left and right derivatives of I_m exist everywhere. When (i), (ii) and (iii) hold it is then known that I_m is an absolutely continuous function. For a proof of (i), (ii) and (iii) we refer the reader to [4].

Let J_m be the restriction of the isoperimetric profile of a half space of \mathbf{R}^n to the interval $(0, \text{vol } E_m)$, and f, g be the inverse functions of I_m, J_m , respectively. We know that

$$g'(a) = \frac{1}{J'_m(g(a))} = \frac{1}{(n-1)H_0(a)},$$

where $H_0(a)$ is the mean curvature of the hemisphere of area a . We also know that, when f' exists,

$$f'(a) = \frac{1}{I'_m(f(a))} = \frac{1}{(n-1)H},$$

where H is the mean curvature in the interior of E_m of any isoperimetric region of volume $f(a)$. From Proposition 4.1 we obtain that $g'(a) \geq f'(a)$ a. e. As f, g are absolutely continuous then $g(a) \geq f(a)$. Since J_m is increasing it easily follows that $I_m \geq J_m$.

If equality holds for some v_0 , then for $a_0 = J_k(v_0) = I_k(v_0)$ we have $g(a_0) = f(a_0)$. Since $g' \geq f'$ we obtain that $f \equiv g$ in the interval $(0, a_0)$ and so $H_0(a_0)^{-1} = H(a_0)^{-1}$. If Ω_0 is any isoperimetric region of volume v_0 then Proposition 4.1 implies that Ω_0 is isometric to a half ball in \mathbf{R}^n of volume v_0 .

Finally let $\Omega \subset \mathbf{R}_C^n$ be relatively compact with smooth boundary. Then $\Omega \subset E_m$, for some m , and

$$\mathcal{P}_C(\Omega) \geq I_m(\text{vol } \Omega) \geq I_{\mathbf{H}^n}(\text{vol } \Omega).$$

If equality holds then Ω is an isoperimetric region in E_m and $I_m(\text{vol } \Omega) = I_{\mathbf{H}^n}(\text{vol } \Omega)$. By the discussion in the above paragraph, Ω is isometric to a half ball in \mathbf{R}^n of volume $\text{vol } \Omega$. \square

Finally we show that the relative isoperimetric inequality (1.1) also holds outside any convex domain in \mathbf{R}^n , with no additional assumptions on the regularity of its boundary

Theorem 5.3. *If $C \subset \mathbf{R}^n$ is any closed convex set with interior points and $D \subset \mathbf{R}_C^n$ is a bounded set with finite perimeter, then*

$$(\text{area}(\partial D)_C)^n \geq \frac{1}{2} n^n \omega_n (\text{vol } D)^{n-1}.$$

Proof. Using standard results on the Hausdorff metric, we can find a sequence of convex sets with smooth boundary $C_m \subset \mathbf{R}^n$, and with $C \subset C_m$ for all $m \in \mathbb{N}$, converging locally in the Hausdorff distance to C . Let $D \subset \mathbf{R}_C^n$ be a bounded set with $(\partial D)_C$ smooth. Define $D_m := D \cap (\mathbf{R}^n)_{C_m}$. Then $\lim_{m \rightarrow \infty} \text{vol } D_m = \text{vol } D$ and $\mathcal{P}_C(D) \geq \mathcal{P}_{C_m}(D_m)$. Since, by Theorem 5.1, the relative isoperimetric inequality (1.1) is satisfied in $(\mathbf{R}^n)_{C_m}$, we have

$$(\text{area}(\partial D)_C)^n \geq (\text{area}(\partial D_m)_{C_m})^n \geq \frac{1}{2} n^n \omega_n (\text{vol } D_m)^{n-1}.$$

Taking limits when $m \rightarrow \infty$, we get (1.1). □

Remark 5.4. Reasoning as in [4], one can easily see that equality is never attained if C is strictly convex. The analysis of equality in the isoperimetric inequality for a general convex set cannot be treated with the tools used in this paper.

REFERENCES

- [1] *T. E. Cecil and P. J. Ryan.*, Tight and taut immersions of manifolds. Pitman (Advanced Publishing Program), Boston, Mass., 1985.
- [2] *J. Choe*, Relative isoperimetric inequality for domains outside a convex set, *Archives Inequalities Appl.* **1**, (2003), 241-250.
- [3] *J. Choe, M. Ghomi and M. Ritoré*, Total positive curvature of hypersurfaces with convex boundary, *J. Differential Geom.*, **72** (2006), 129–147.
- [4] *J. Choe and M. Ritoré*, The relative isoperimetric inequality in Cartan-Hadamard 3-manifolds, preprint, 2004.
- [5] *H. Federer*, Geometric Measure Theory, Die Grundlehren der mathematischen Wissenschaften, Band 153, Springer-Verlag New York Inc., New York, 1969.
- [6] *E. Giusti*, Minimal surfaces and functions of bounded variation, *Monographs in Mathematics*, no. 80, Birkhäuser Verlag, Basel-Boston, 1984.
- [7] *E. Gonzalez, U. Massari and I. Tamanini*, On the regularity of boundaries of sets minimizing perimeter with a volume constraint, *Indiana Univ. Math. J.* **32** (1983), no. 1, 25–37
- [8] *M. Grüter*, Optimal regularity for codimension one minimal surfaces with a free boundary. *Manusc. Math.* **58** (1987), 295–343
- [9] *M. Grüter*, Boundary regularity for solutions of a partitioning problem. *Arch. Rational Mech. Anal.* **97** (1987), no. 3, 261–270.

- [10] *Wu-Yi Hsiang*, On soap bubbles and isoperimetric regions in noncompact symmetric spaces. I, *Tohoku Math. J. (2)* **44** (1992), no. 2, 151–175.
- [11] *I. Kim*, An optimal relative isoperimetric inequality in concave cylindrical domains in \mathbf{R}^n , *J. Inequalities Appl.* **1** (2000), 97-102.
- [12] *P. Li, S. T. Yau*, A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces, *Invent. Math.* **69** (1982) no. 2, 269–291.
- [13] *M. Ritoré and C. Rosales*, Existence and characterization of regions minimizing perimeter under a volume constraint inside Euclidean cones, *Trans. Amer. Math. Soc.* **356** (2004), no. 11, 4601–4622.
- [14] *R. Schneider*, *Convex bodies: the Brunn-Minkowski theory*. Cambridge University Press, Cambridge, 1993.
- [15] *E. Stredulinsky and W. P. Ziemer*, Area minimizing sets subject to a volume constraint in a convex set, *J. Geom. Anal.* **7** (1997), no. 4, 653–677.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL, 151-742, KOREA

E-mail address: `choe@math.snu.ac.kr`

URL: `www.math.snu.ac.kr/~choe`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332

E-mail address: `ghomi@math.gatech.edu`

URL: `www.math.gatech.edu/~ghomi`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, E-18071 GRANADA, ESPAÑA

E-mail address: `ritore@ugr.es`

URL: `www.ugr.es/~ritore`