

## New study highlights the importance of physical activity during the early weeks of pregnancy

10/01/2020

Research news

**Researchers at the University of Granada (UGR) have analysed for the first time the link between sedentary time vs. objectively-measured physical activity levels and systemic concentrations of glycaemic, lipid, and inflammatory markers among pregnant women**

Pregnancy is a critical stage characterised by complex changes in endocrine and immunometabolic levels, during which the woman's physiology is modulated to sustain her health and that of the baby. However, a malfunction in these mechanisms during the early stages of pregnancy may subsequently lead to complications and adverse consequences for foetal development and the future health of both mother and child.

In view of this issue, in recent years a new field of research has evolved, focusing on the quest for strategies that enable pregnant women to better regulate and maintain these mechanisms during pregnancy, thus facilitating pregnancies with fewer complications.

Given the proven influence of lifestyle on the intrauterine environment, reducing the time spent by the mother on sedentary behaviours and increasing her physical activity levels may constitute alternative-complementary tools for controlling and regulating these immunometabolic responses and therefore promoting healthier pregnancies. However, only a few studies have been conducted to date dealing with women during the early stages of pregnancy, and the results are ambiguous. Furthermore, this issue has never been explored in pregnant women with no severe



metabolic disruptions.

Researchers from the University of Granada (UGR) have recently published a study in the Scandinavian Journal of Medicine and Science in Sports, in which they analyse for the first time the link between sedentary time vs. objectively-measured physical activity levels and systemic concentrations of glycaemic, lipid, and inflammatory markers among women in the early stages of pregnancy (n= 50 pregnant women; 33 years of age; body mass index 24.2kg/m<sup>2</sup>; gestational stage 17 weeks). This study is part of the GESTAFIT (“Gestation and Fitness”) project, which was carried out at the UGR during the period 2015–2018.

The authors of this study observed that higher levels of physical activity at a moderate-vigorous intensity and complying with the specific physical activity recommendations for this population during the first weeks of pregnancy were associated with lower concentrations of interleukin 1 $\beta$  and interferon- $\gamma$  and higher levels of interleukin 8. That is, physical activity was linked to physiological markers related to inflammation, angiogenesis, metabolic regulation, and so on. However, the researchers did not identify any relationship between sedentary time or physical activity levels and glycaemic and lipid markers.

They concluded that spending more time conducting physical activity of moderate-vigorous intensity may help modulate these immunometabolic responses during the early stages of pregnancy in ‘healthy’ pregnant women. Given the importance of maintaining an adequate balance of these mechanisms to promote more successful pregnancies, the results of the present study could be relevant to the clinical setting.

Pedro Acosta-Manzano and Francisco M. Acosta, the main authors of the study, explain: “If future intervention studies confirm these results in larger samples of pregnant women, different healthcare institutions might consider implementing strategies aimed at increasing physical activity levels (mainly of moderate-vigorous intensity) as a complementary tool to control the balance between these immunometabolic responses in pregnant women with no severe metabolic disruptions.”

### **Bibliography:**

Acosta-Manzano, P., Acosta, F.M., Femia, P., Coll-Risco, I., Segura-Jiménez, V., Diaz-Castro, J., Ochoa-Herrera J., Van Poppel, M.N.M., and Aparicio, V.A., 'Association of sedentary time and physical activity levels with glycaemic , lipid and inflammatory markers in early pregnancy: The GESTAFIT Project', Scandinavian Journal of Medicine and Science in Sports. <https://doi.org/10.1111/sms.13547>



*Part of the GESTAFIT team*

**Media enquiries:**

Pedro Acosta Manzano

Department of Physical and Sports Education, University of Granada

Tel.: +34 958 244393

Email: [acostapedro23@ugr.es](mailto:acostapedro23@ugr.es)