

# The group of isometries of a Banach space and duality

**Miguel Martín**

<http://www.ugr.es/local/mmartins>



February 28th, 2008 – Universidad de Murcia

# Outline

- 1 Notation and objective
- 2 The tool: numerical range of operators
  - Definitions
  - Relationship with semigroups of operators
- 3 The example
- 4 Some related results
  - Finite-dimensional spaces
  - Numerical index and duality

# Notation and objective

## Basic notation

$X$  Banach space over  $\mathbb{K}$  ( $= \mathbb{R}$  or  $\mathbb{C}$ ).

- $S_X$  unit sphere,  $B_X$  unit ball,
- $X^*$  dual space,
- $L(X)$  bounded linear operators,
- $\text{Iso}(X)$  surjective linear isometries,
- $T^* \in L(X^*)$  adjoint operator of  $T \in L(X)$ .

## Main Objective

We **construct** a real Banach space  $X$  such that

- $\text{Iso}(X)$  does not contain uniformly continuous one-parameter semigroups.
- But  $\text{Iso}(X^*)$  contains infinitely many uniformly continuous one-parameter semigroups.

## The tool: numerical range of operators



F. F. Bonsall and J. Duncan

*Numerical Ranges of Operators on Normed Spaces and of Elements of Normed Algebras.*

London Math. Soc. Lecture Note Series, 1971.



F. F. Bonsall and J. Duncan

*Numerical Ranges II.*

London Math. Soc. Lecture Note Series, 1973.



H. P. Rosenthal

The Lie algebra of a Banach space.

in: *Banach spaces* (Columbia, Mo., 1984), LNM, Springer, 1985.

# Hilbert spaces

## Hilbert space Numerical range (Toeplitz, 1918)

- $A$   $n \times n$  real or complex matrix

$$W(A) = \{(Ax \mid x) : x \in \mathbb{K}^n, (x \mid x) = 1\}.$$

- $H$  real or complex Hilbert space,  $T \in L(H)$ ,

$$W(T) = \{(Tx \mid x) : x \in H, \|x\| = 1\}.$$

## Some properties

$H$  Hilbert space,  $T \in L(H)$ :

- $W(T)$  is convex.
- In the complex case,  $\overline{W(T)}$  contains the spectrum of  $T$ .
- If, moreover,  $T$  is normal,  $\overline{W(T)} = \overline{\text{co}} \text{Sp}(T)$ .

# Banach spaces

## Banach space numerical range (Bauer 1962; Lumer, 1961)

$X$  Banach space,  $T \in L(X)$ ,

$$V(T) = \{x^*(Tx) : x^* \in S_{X^*}, x \in S_X, x^*(x) = 1\}$$

## Some properties

$X$  Banach space,  $T \in L(X)$ :

- $V(T)$  is connected (not necessarily convex).
- In the complex case,  $\overline{V(T)}$  contains the spectrum of  $T$ .
- Actually,

$$\overline{\text{co}} \text{Sp}(T) = \bigcap \overline{\text{co}} V(T),$$

the intersection taken over all numerical ranges  $V(T)$  corresponding to equivalent norms on  $X$ .



## Numerical radius

$X$  real or complex Banach space,  $T \in L(X)$ ,

$$v(T) = \sup \{ |\lambda| : \lambda \in V(T) \}.$$

- $v$  is a seminorm with  $v(T) \leq \|T\|$ .
- $v(T) = v(T^*)$  for every  $T \in L(X)$ .

## Numerical index (Lumer, 1968)

$X$  real or complex Banach space,

$$\begin{aligned} n(X) &= \inf \{ v(T) : T \in L(X), \|T\| = 1 \} \\ &= \max \{ k \geq 0 : k\|T\| \leq v(T) \forall T \in L(X) \}. \end{aligned}$$

## Remarks

- $n(X) = 1$  iff  $v(T) = \|T\|$  for every  $T \in L(X)$ .
- If there is  $T \neq 0$  with  $v(T) = 0$ , then  $n(X) = 0$ .
- The converse is not true.

## Relationship with semigroups of operators

### A motivating example

A real or complex  $n \times n$  matrix. TFAE:

- $A$  is skew-adjoint (i.e.  $A^* = -A$ ).
- $\operatorname{Re}(Ax \mid x) = 0$  for every  $x \in H$ .
- $B = \exp(\rho A)$  is unitary for every  $\rho \in \mathbb{R}$  (i.e.  $B^*B = \operatorname{Id}$ ).

### In term of Hilbert spaces

$H$  ( $n$ -dimensional) Hilbert space,  $T \in L(H)$ . TFAE:

- $\operatorname{Re} W(T) = \{0\}$ .
- $\exp(\rho T) \in \operatorname{Iso}(H)$  for every  $\rho \in \mathbb{R}$ .

### For general Banach spaces

$X$  Banach space,  $T \in L(X)$ . TFAE:

- $\operatorname{Re} V(T) = \{0\}$ .
- $\exp(\rho T) \in \operatorname{Iso}(X)$  for every  $\rho \in \mathbb{R}$ .



# Characterizing uniformly continuous semigroups of operators

## Theorem

$X$  real or complex Banach space,  $T \in L(X)$ . TFAE:

- $\operatorname{Re} V(T) = \{0\}$ .
- $\|\exp(\rho T)\| \leq 1$  for every  $\rho \in \mathbb{R}$ .
- $\{\exp(\rho T) : \rho \in \mathbb{R}_0^+\} \subset \operatorname{Iso}(X)$ .
- $T$  belongs to the tangent space of  $\operatorname{Iso}(X)$  at  $\operatorname{Id}$ , i.e. exists a function  $f : [-1, 1] \rightarrow \operatorname{Iso}(X)$  with  $f(0) = \operatorname{Id}$  and  $f'(0) = T$ .
- $\lim_{\rho \rightarrow 0} \frac{\|\operatorname{Id} + \rho T\| - 1}{\rho} = 0$ , i.e. the derivative or the norm of  $L(X)$  at  $\operatorname{Id}$  in the direction of  $T$  is null.

## Main consequence for us

If  $X$  is a **real** Banach space with  $n(X) > 0$ , then  $\operatorname{Iso}(X)$  is “small”:

- it does not contain any uniformly continuous one-parameter semigroups,
- the tangent space of  $\operatorname{Iso}(X)$  at  $\operatorname{Id}$  is zero.

## The example



M. Martín

The group of isometries of a Banach space and duality.  
*preprint.*

## The main example

### The construction

$E$  separable Banach space. We construct a Banach space  $X(E)$  such that

$$n(X(E)) = 1 \quad \text{and} \quad X(E)^* \cong E^* \oplus_1 L_1(\mu)$$

### The main consequence

Take  $E = \ell_2$  (real). Then

- $n(X(\ell_2)) = 1$ , so  $\text{Iso}(X(\ell_2))$  is “small”.
- Since  $X(\ell_2)^* \cong \ell_2 \oplus_1 L_1(\mu)$ , given  $S \in \text{Iso}(\ell_2)$ , the operator

$$T = \begin{pmatrix} S & 0 \\ 0 & \text{Id} \end{pmatrix}$$

is a surjective isometry of  $X(\ell_2)^*$ .

- Therefore,  $\text{Iso}(X(\ell_2)^*)$  contains infinitely many semigroups of isometries.

## Sketch of the construction I

Define (viewing  $E \hookrightarrow C[0, 1]$ )

$$Y = \{f \in C([0, 1] \times [0, 1]) : f(\cdot, 0) = 0\}$$

$$X(E) = \{f \in C([0, 1] \times [0, 1]) : f(\cdot, 0) \in E\}$$

We need

$$X(E)^* \cong E^* \oplus_1 L_1(\mu) \quad \&$$

$$n(X(E)) = 1$$

Proving that  $X(E)^* \cong E^* \oplus_1 L_1(\mu)$ 

- $Y$  is an  $M$ -ideal of  $C([0, 1] \times [0, 1])$ , so  $Y$  is an  $M$ -ideal of  $X(E)$ .
- This means that  $X(E)^* \cong Y^\perp \oplus_1 Y^*$ .
- $Y^* \cong L_1(\mu)$  for some measure  $\mu$ ;  $Y^\perp \cong (X(E)/Y)^*$ .
- Define  $\Phi : X(E) \rightarrow E$  by  $\Phi(f) = f(\cdot, 0)$ .
  - $\|\Phi\| \leq 1$  and  $\ker \Phi = Y$ .
  - $\tilde{\Phi} : X(E)/Y \rightarrow E$  is a surjective isometry since:
    - $\{g \in E : \|g\| < 1\} \subseteq \Phi(\{f \in X(E) : \|f\| < 1\})$ .
- Therefore,  $Y^\perp \cong (X(E)/Y)^* \cong E^*$ .

# Sketch of the construction II

Define (viewing  $E \hookrightarrow C[0, 1]$ )

$$Y = \{f \in C([0, 1] \times [0, 1]) : f(\cdot, 0) = 0\}$$

$$X(E) = \{f \in C([0, 1] \times [0, 1]) : f(\cdot, 0) \in E\}$$

We need

$$X(E)^* \cong E^* \oplus L_1(\mu) \quad \& \quad n(X(E)) = 1$$

Proving that  $n(X(E)) = 1$

- Fix  $T \in L(X(E))$ . Find  $f_0 \in X(E)$  and  $\xi_0 \in ]0, 1] \times [0, 1]$  such that  $|[Tf_0](\xi_0)| \sim \|T\|$ .
- Consider the non-empty open set

$$V = \{\xi \in ]0, 1] \times [0, 1] : f_0(\xi) \sim f_0(\xi_0)\}$$

and find  $\varphi : [0, 1] \times [0, 1] \rightarrow [0, 1]$  continuous with  $\text{supp}(\varphi) \subset V$  and  $\varphi(\xi_0) = 1$ .

- Write  $f_0(\xi_0) = \lambda\omega_1 + (1 - \lambda)\omega_2$  with  $|\omega_i| = 1$ , and consider the functions

$$f_i = (1 - \varphi)f_0 + \varphi\omega_i \text{ for } i = 1, 2.$$

- Then,  $f_i \in Y \subset X(E)$ ,  $\|f_i\| \leq 1$ , and

$$\|f_0 - (\lambda f_1 + (1 - \lambda)f_2)\| = \|\varphi f_0 - \varphi f_0(\xi_0)\| \sim 0.$$

- Therefore, there is  $i \in \{1, 2\}$  such that  $|[T(f_i)](\xi_0)| \sim \|T\|$ , but now  $|f_i(\xi_0)| = 1$ .
- Equivalently,

$$|\delta_{\xi_0}(T(f_i))| \sim \|T\| \quad \text{and} \quad |\delta_{\xi_0}(f_i)| = 1,$$

meaning that  $v(T) \sim \|T\|$ .

## Some related results



K. Boyko, V. Kadets, M. Martín, and D. Werner.  
Numerical index of Banach spaces and duality.  
*Math. Proc. Cambridge Philos. Soc.* (2007).



M. Martín, J. Merí, and A. Rodríguez-Palacios.  
Finite-dimensional spaces with numerical index zero.  
*Indiana U. Math. J.* (2004).



H. P. Rosenthal  
The Lie algebra of a Banach space.  
in: *Banach spaces* (Columbia, Mo., 1984), LNM, Springer, 1985.

# Isometries in finite-dimensional spaces

## Theorem

Let  $X$  be a finite-dimensional **real** space. TFAE:

- $\text{Iso}(X)$  is infinite.
- $n(X) = 0$ .
- There is  $T \in L(X)$ ,  $T \neq 0$ , with  $v(T) = 0$ .

## Examples of spaces of this kind

- 1 Hilbert spaces.
- 2  $X_{\mathbb{R}}$ , the real space subjacent to any complex space  $X$ .
- 3 An absolute sum of any real space and one of the above.
- 4 Moreover, if  $X = X_0 \oplus X_1$  where  $X_1$  is complex and

$$\|x_0 + e^{i\theta} x_1\| = \|x_0 + x_1\| \quad (x_0 \in X_0, x_1 \in X_1, \theta \in \mathbb{R}).$$

(Note that the other 3 cases are included here)

## Question

Can every Banach space  $X$  with  $n(X) = 0$  be decomposed as in 4 ?

## Negative answer I

### Infinite-dimensional case

There is an infinite-dimensional real Banach space  $X$  with  $n(X) = 0$  but  $X$  is polyhedral. In particular,  $X$  does not contain  $\mathbb{C}$  isometrically.

### The example is

$$X = \left[ \bigoplus_{n \geq 2} X_n \right]_{c_0}$$

$X_n$  is the two-dimensional space whose unit ball is the regular polygon of  $2n$  vertices.

### Note

Such an example is not possible in the finite-dimensional case.



## (Quasi affirmative) negative answer II

## Finite-dimensional case

$X$  finite-dimensional real space. TFAE:

- $n(X) = 0$ .
- $X = X_0 \oplus X_1 \oplus \cdots \oplus X_n$  such that
  - $X_0$  is a (possible null) real space,
  - $X_1, \dots, X_n$  are non-null complex spaces,

there are  $\rho_1, \dots, \rho_n$  **rational** numbers, such that

$$\left\| x_0 + e^{i\rho_1\theta} x_1 + \cdots + e^{i\rho_n\theta} x_n \right\| = \left\| x_0 + x_1 + \cdots + x_n \right\|$$

for every  $x_i \in X_i$  and every  $\theta \in \mathbb{R}$ .

## Example

$$X = (\mathbb{R}^4, \|\cdot\|), \|(a, b, c, d)\| = \frac{1}{4} \int_0^{2\pi} \left| \operatorname{Re} \left( e^{2it}(a + ib) + e^{it}(c + id) \right) \right| dt.$$

Then  $n(X) = 0$  but the unique possible decomposition is  $X = \mathbb{C} \oplus \mathbb{C}$  with

$$\left\| e^{it} x_1 + e^{2it} x_2 \right\| = \|x_1 + x_2\|.$$

# The Lie-algebra of a Banach space

## Lie-algebra

$X$  real Banach space,  $\mathcal{Z}(X) = \{T \in L(X) : v(T) = 0\}$ .

- When  $X$  is finite-dimensional,  $\text{Iso}(X)$  is a Lie-group and  $\mathcal{Z}(X)$  is the tangent space (i.e. its Lie-algebra).

## Remark

If  $\dim(X) = n$ , then

$$0 \leq \dim(\mathcal{Z}(X)) \leq \frac{n(n-1)}{2}.$$

## An open problem

Given  $n \geq 3$ , which are the possible  $\dim(\mathcal{Z}(X))$  over all  $n$ -dimensional  $X$ 's?

## Observation (Javier Merí, PhD)

When  $\dim(X) = 3$ ,  $\dim(\mathcal{Z}(X))$  cannot be 2.

# Numerical index of Banach spaces

## Numerical index (Lumer, 1968)

$X$  real or complex Banach space,

$$n(X) = \inf \{v(T) : T \in L(X), \|T\| = 1\} = \max \{k \geq 0 : k\|T\| \leq v(T) \forall T \in L(X)\}.$$

## Some examples

①  $C(K)$ ,  $L_1(\mu)$  have numerical index 1.

②  $H$  Hilbert space,  $\dim(H) > 1$ , then

$$n(H) = 0 \text{ real case} \quad n(H) = \frac{1}{2} \text{ complex case.}$$

③  $n(L_p[0, 1]) = n(\ell_p)$  but both are unknown.

④ If  $X_n$  is the two-dimensional space such that  $B_{X_n}$  is a  $2n$ -polygon, then

$$n(X_n) = \tan\left(\frac{\pi}{2n}\right) \text{ if } n \text{ is even} \quad n(X_n) = \sin\left(\frac{\pi}{2n}\right) \text{ if } n \text{ is odd.}$$

⑤ If  $X$  is a  $C^*$ -algebra or the predual of a von Neumann algebra, then  $n(X) = 1$  if the algebra is commutative and  $n(X) = 1/2$  otherwise.

# Numerical index and duality

## Proposition

$X$  Banach space.

- $v(T^*) = v(T)$  for every  $T \in L(X)$ .
- Therefore,  $n(X^*) \leq n(X)$ .

## Question

Is it always  $n(X) = n(X^*)$  ?

## Another example

- It is known: if  $X$  or  $X^*$  is a  $C^*$ -algebra, then  $n(X) = n(X^*)$ .
- Consider  $Y = X(K(\ell_2))$ . Then

$$n(Y) = 1 \quad \text{and} \quad Y^* \cong K(\ell_2)^* \oplus_1 L_1(\mu).$$

Then,  $Y^{**} \cong L(\ell_2) \oplus_\infty L_\infty(\mu)$  is a  $C^*$ -algebra but  $n(Y^*) \leq n(K(\ell_2)) = 1/2$ .

# Numerical index and duality

## Remark

In the example  $n(X(\ell_2)) > n(X(\ell_2)^*)$ , one finds that  $X(\ell_2)^*$  has another predual (namely,  $\ell_2 \oplus_\infty Y$ ) for which the numerical index coincides with the numerical index of its dual.

## Open problems

We look for sufficient conditions assuring the equality between the numerical index of a Banach space and the one of its dual.

- ① Asplundness is not such a property.
- ② What's about RNP ?
- ③ What's about if  $X^*$  has a unique predual ?
- ④ What's about if  $X$  does not contains a copy of  $c_0$  ?

## Theorem

If  $X$  is a separable Banach space containing (an isomorphic copy of)  $c_0$ , then there is an equivalent norm  $|\cdot|$  on  $X$  such that

$$n((X, |\cdot|)^*) < n((X, |\cdot|)).$$

# The group of isometries of a Banach space and duality

**Miguel Martín**

<http://www.ugr.es/local/mmartins>



February 28th, 2008 – Universidad de Murcia