

Isometries of Banach spaces and duality

Miguel Martín

<http://www.ugr.es/local/mmartins>



ugr

Universidad
de Granada



Summer Conference on General Topology and its Applications

New York, July 2011

Basic notation and main objective

Notation

X Banach space.

- S_X unit sphere, B_X closed unit ball.
- X^* dual space.
- $L(X)$ bounded linear operators.
- $W(X)$ weakly compact linear operators.
- $\text{Iso}(X)$ surjective isometries group.

Objective

★ Construct a Banach space X with “small” $\text{Iso}(X)$ and “big” $\text{Iso}(X^*)$.

A previous attempt

M., 2008

There is X such that

- $\text{Iso}(X)$ does not contain uniformly continuous semigroups of isometries;
- $\text{Iso}(X^*) \supset \text{Iso}(\ell_2)$ and, therefore, $\text{Iso}(X^*)$ contains infinitely many uniformly continuous semigroups of isometries.
- But $\text{Iso}(X)$ contains infinitely many **strongly continuous** semigroups of isometries.

Question we are going to solve

Is it possible to produce a space X such that $\text{Iso}(X^*) \supset \text{Iso}(\ell_2)$ but $\text{Iso}(X)$ is “smaller” (for instance, it does not contain strongly continuous semigroups) ?

Motivation

X Banach space.

Autonomous dynamic system

$$(\diamond) \quad \begin{cases} x'(t) = Ax(t) \\ x(0) = x_0 \end{cases} \quad x_0 \in X, A \text{ linear, closed, densely defined.}$$

One-parameter semigroup of operators

$\Phi : \mathbb{R}_0^+ \rightarrow L(X)$ such that $\Phi(t+s) = \Phi(t)\Phi(s) \forall t, s \in \mathbb{R}_0^+, \Phi(0) = \text{Id}$.

- *Uniformly continuous*: $\Phi : \mathbb{R}_0^+ \rightarrow (L(X), \|\cdot\|)$ continuous.
- *Strongly continuous*: $\Phi : \mathbb{R}_0^+ \rightarrow (L(X), \text{SOT})$ continuous.

Relationship (Hille-Yoshida, 1950's)

- *Bounded case*:
 - If $A \in L(X) \implies \Phi(t) = \exp(tA)$ solution of (\diamond) uniformly continuous.
 - Φ uniformly continuous $\implies A = \Phi'(0) \in L(X)$ and Φ solution of (\diamond) .
- *Unbounded case*:
 - Φ strongly continuous $\implies A = \Phi'(0)$ closed and Φ solution of (\diamond) .
 - If (\diamond) has solution Φ strongly continuous $\implies A = \Phi'(0)$ and $\Phi(t) = \text{"exp}(tA)\text{"}$.

What we are going to show

The example

we will construct X such that

$$\text{Iso}(X) = \{\pm \text{Id}\} \quad \text{but} \quad \text{Iso}(X^*) \supset \text{Iso}(\ell_2).$$

The tools

- **Extremely non-complex Banach spaces:** spaces X such that $\|\text{Id} + T^2\| = 1 + \|T^2\|$ for every $T \in L(X)$.
- **Koszmider type compact spaces:** topological compact spaces K such that $C(K)$ has few operators.

The talk is based on the papers



P. Koszmider, M. Martín, and J. Merí.
Extremely non-complex $C(K)$ spaces.
J. Math. Anal. Appl. (2009).



P. Koszmider, M. Martín, and J. Merí.
Isometries on extremely non-complex Banach spaces.
J. Inst. Math. Jussieu (2011).



M. Martín
The group of isometries of a Banach space and duality.
J. Funct. Anal. (2008).

Sketch of the talk

- 1 Introduction
- 2 Extremely non-complex Banach spaces: motivation and examples
- 3 Isometries on extremely non-complex spaces

Extremely non-complex Banach spaces: motivation and examples

- 1 Introduction
- 2 Extremely non-complex Banach spaces: motivation and examples
 - Complex structures
 - The first examples: $C(K)$ spaces with few operators
 - More $C(K)$ -type examples
 - Further examples
- 3 Isometries on extremely non-complex spaces

Complex structures

Definition

X has **complex structure** if there is $T \in L(X)$ such that $T^2 = -\text{Id}$.

Some remarks

- This gives a structure of vector space over \mathbb{C} :

$$(\alpha + i\beta)x = \alpha x + \beta T(x) \quad (\alpha + i\beta \in \mathbb{C}, x \in X)$$

- Defining

$$\|x\| = \max\{\|e^{i\theta}x\| : \theta \in [0, 2\pi]\} \quad (x \in X)$$

one gets that $(X, \|\cdot\|)$ is a complex Banach space.

- If T is an isometry, then actually the given norm of X is complex.
- Conversely, if X is a complex Banach space, then

$$T(x) = ix \quad (x \in X)$$

satisfies $T^2 = -\text{Id}$ and T is an isometry.

Complex structures II

Some examples

- 1 If $\dim(X) < \infty$, X has complex structure iff $\dim(X)$ is even.
- 2 If $X \simeq Z \oplus Z$ (in particular, $X \simeq X^2$), then X has complex structure.
- 3 There are infinite-dimensional Banach spaces without complex structure:
 - **Dieudonné, 1952:** the James' space \mathcal{J} (since $\mathcal{J}^{**} \equiv \mathcal{J} \oplus \mathbb{R}$).
 - **Szarek, 1986:** uniformly convex examples.
 - **Gowers-Maurey, 1993:** their H.I. space.

Definition

X is **extremely non-complex** if $\text{dist}(T^2, -\text{Id})$ is the maximum possible, i.e.

$$\|\text{Id} + T^2\| = 1 + \|T^2\| \quad (T \in L(X))$$

Question (Gilles Godefroy, private communication, 2005)

Is there any $X \neq \mathbb{R}$ such that $\|\text{Id} + T^2\| = 1 + \|T^2\|$ for every $T \in L(X)$?

Weak multipliers

Weak multipliers

Let K be a compact space. $T \in L(C(K))$ is a **weak multiplier** if

$$T^* = g\text{Id} + S$$

where g is a Borel function and S is weakly compact.

Theorem

K perfect, $T \in L(C(K))$ weak multiplier $\implies \|\text{Id} + T^2\| = 1 + \|T^2\|$

Examples (Koszmider, 2004)

There are infinitely many different perfect compact spaces K such that all operators on $C(K)$ are weak multipliers.

They are called **weak Koszmider spaces**.

Corollary

There are infinitely many non-isomorphic extremely non-complex spaces.

More $C(K)$ -type examples

More $C(K)$ type examples

There are perfect compact spaces K_1, K_2 such that:

- $C(K_1)$ and $C(K_2)$ are extremely non-complex,
- $C(K_1)$ contains a complemented copy of $C(\Delta)$.
- $C(K_2)$ contains a (1-complemented) isometric copy of ℓ_∞ .

Observation

- $C(K_1)$ and $C(K_2)$ have operators which are not weak multipliers.
- They are not indecomposable spaces.

Further examples

Spaces $C_E(K\|L)$

K compact, $L \subset K$ closed nowhere dense, $E \subset C(L)$. Define

$$C_E(K\|L) := \{f \in C(K) : f|_L \in E\}.$$

Observation

$C_0(K\|L)$ is an M -ideal in $C_E(K\|L)$, meaning that

$$C_E(K\|L)^* \cong E^* \oplus_1 C_0(K\|L)^*$$

Theorem

K perfect weak Koszmider, L closed nowhere dense, $E \subset C(L)$

$\implies C_E(K\|L)$ is extremely non-complex.

Isometries on extremely non-complex spaces

- 1 Introduction
- 2 Extremely non-complex Banach spaces: motivation and examples
- 3 **Isometries on extremely non-complex spaces**
 - Isometries on extremely non-complex spaces
 - Isometries on extremely non-complex $C_E(K||L)$ spaces
 - The main example

Isometries on extremely non-complex spaces

Theorem

X extremely non-complex.

- $T \in \text{Iso}(X) \implies T^2 = \text{Id}$.
- $T_1, T_2 \in \text{Iso}(X) \implies T_1 T_2 = T_2 T_1$.
- $T_1, T_2 \in \text{Iso}(X) \implies \|T_1 - T_2\| \in \{0, 2\}$.
- $\Phi : \mathbb{R}_0^+ \longrightarrow \text{Iso}(X)$ one-parameter semigroup $\implies \Phi(\mathbb{R}_0^+) = \{\text{Id}\}$.

Consequences

- $\text{Iso}(X)$ is a Boolean group for the composition operation.
- $\text{Iso}(X)$ identifies with the set $\text{Unc}(X)$ of unconditional projections on X :

$$\begin{aligned} P \in \text{Unc}(X) &\iff P^2 = P, 2P - \text{Id} \in \text{Iso}(X) \\ &\iff P = \frac{1}{2}(\text{Id} - T), T \in \text{Iso}(X), T^2 = \text{Id}. \end{aligned}$$

Extremely non-complex $C_E(K\|L)$ spaces.

Remember

K perfect weak Koszmider, L closed nowhere dense, $E \subset C(L)$
 $\implies C_E(K\|L)$ is extremely non-complex and $C_E(K\|L)^* \cong E^* \oplus_1 C_0(K\|L)^*$.

Proposition

K perfect $\implies \exists L \subset K$ closed nowhere dense with $C[0,1] \subset C(L)$.

A good example

Take K perfect weak Koszmider, $L \subset K$ closed nowhere dense with
 $E = \ell_2 \subset C[0,1] \subset C(L)$:

- $C_{\ell_2}(K\|L)$ has no non-trivial one-parameter semigroup of isometries.
- $C_{\ell_2}(K\|L)^* \cong \ell_2 \oplus_1 C_0(K\|L)^* \implies \text{Iso}(C_{\ell_2}(K\|L)^*) \supset \text{Iso}(\ell_2)$.

But we are able to give a better result...

Isometries on extremely non-complex $C_E(K||L)$ spaces

Theorem

$C_E(K||L)$ extremely non-complex, $T \in \text{Iso}(C_E(K||L))$
 \implies exists $\theta : K \setminus L \rightarrow \{-1, 1\}$ continuous such that

$$[T(f)](x) = \theta(x)f(x) \quad (x \in K \setminus L, f \in C_E(K||L))$$

Consequences: cases $E = C(L)$ and $E = 0$

- $C(K)$ extremely non-complex, $\varphi : K \rightarrow K$ homeomorphism $\implies \varphi = \text{id}$
- $C_0(K \setminus L) \cong C_0(K||L)$ extremely non-complex, $\varphi : K \setminus L \rightarrow K \setminus L$ homeomorphism $\implies \varphi = \text{id}$

Consequence: connected case

If K and $K \setminus L$ are connected, then

$$\text{Iso}(C_E(K||L)) = \{-\text{Id}, +\text{Id}\}$$

The main example

Koszmider, 2004

$\exists \mathcal{K}$ connected weak Koszmider space such that $\mathcal{K} \setminus F$ is connected if $|F| < \infty$.

Important observation on the construction above

There is $\mathcal{L} \subset \mathcal{K}$ closed and nowhere dense, with

- $\mathcal{K} \setminus \mathcal{L}$ connected
- $C[0,1] \subseteq C(\mathcal{L})$

Consequence: the best example

Consider $X = C_{\ell_2}(\mathcal{K}||\mathcal{L})$. Then:

$$\text{Iso}(X) = \{-\text{Id}, +\text{Id}\} \quad \text{and} \quad \text{Iso}(X^*) \supset \text{Iso}(\ell_2)$$

Proof.

- \mathcal{K} weak Koszmider, \mathcal{L} nowhere dense, $\ell_2 \subset C[0,1] \subset C(\mathcal{L})$
 $\implies X$ well-defined and extremely non-complex.
- $\mathcal{K} \setminus \mathcal{L}$ connected $\implies \text{Iso}(X) = \{-\text{Id}, +\text{Id}\}$.
- $X^* \equiv \ell_2 \oplus_1 C_0(\mathcal{K}||\mathcal{L})^* \implies \text{Iso}(\ell_2) \subset \text{Iso}(X^*)$.