



# 수학교육과 강연회



동국대학교 사범대학 수학교육과

## 강연회 알림

대상 : 수학교육과 학생 및 동국인

일시 : 2012년 11월 8일 4시

장소 : 학림관 401호

문의 : 수학교육과 학과사무실(2260-8749)

제목 : Isometries of  
Banach spaces and  
duality.

연사 : Miguel Martin

University of Granada,  
Spain.

# *Introduction*

---

## Section 1

- 1 Introduction
  - Notation and objective
  - A previous attempt
  - Motivation

## Basic notation and main objective

### Notation

$X$  Banach space.

- $S_X = \{x \in X : \|x\| = 1\}$  unit sphere,  $B_X = \{x \in X : \|x\| \leq 1\}$  closed unit ball.
- $X^*$  dual space.
- $L(X)$  bounded linear operators.
- $W(X)$  weakly compact linear operators.
- $\text{Iso}(X)$  surjective (linear) isometries group.

### Note

- $T : X \rightarrow X$  **isometry**  $\stackrel{\text{def}}{\iff} \|Tx - Ty\| = \|x - y\|$  for all  $x, y \in X$ .
- **(Mazur-Ulam)**  $T$  surjective isometry,  $T(0) = 0 \implies T$  linear.

### Objective

- ★ Construct a Banach space  $X$  with “small”  $\text{Iso}(X)$  and “big”  $\text{Iso}(X^*)$ .

## A previous attempt

M., 2008

There is  $X$  such that

- $\text{Iso}(X)$  does not contain uniformly continuous semigroups of isometries;
- $\text{Iso}(X^*) \supset \text{Iso}(\ell_2)$  and, therefore,  $\text{Iso}(X^*)$  contains infinitely many uniformly continuous semigroups of isometries.
- But  $\text{Iso}(X)$  contains infinitely many **strongly continuous** semigroups of isometries.

Question we are going to solve

Is it possible to produce a space  $X$  such that  $\text{Iso}(X^*) \supset \text{Iso}(\ell_2)$  but  $\text{Iso}(X)$  is “small” (for instance, it does not contain strongly continuous semigroups) ?

# Motivation

$X$  Banach space.

## Autonomous linear dynamical system

$$(\diamond) \quad \begin{cases} x'(t) = Ax(t) \\ x(0) = x_0 \end{cases} \quad x_0 \in X, A : X \rightarrow X \text{ linear, closed, densely defined.}$$

## One-parameter semigroup of operators

$\Phi : \mathbb{R}_0^+ \rightarrow L(X)$  such that  $\Phi(t+s) = \Phi(t)\Phi(s) \forall t, s \in \mathbb{R}_0^+, \Phi(0) = \text{Id}$ .

- *Uniformly continuous*:  $\Phi : \mathbb{R}_0^+ \rightarrow (L(X), \|\cdot\|)$  continuous.
- *Strongly continuous*:  $\Phi : \mathbb{R}_0^+ \rightarrow (L(X), \text{SOT})$  continuous.

## Relationship (Hille-Yoshida, 1950's)

- *Bounded case*:
  - If  $A \in L(X) \implies \Phi(t) = \exp(tA)$  solution of  $(\diamond)$  uniformly continuous.
  - $\Phi$  uniformly continuous  $\implies A = \Phi'(0) \in L(X)$  and  $\Phi$  solution of  $(\diamond)$ .
- *Unbounded case*:
  - $\Phi$  strongly continuous  $\implies A = \Phi'(0)$  closed and  $\Phi$  solution of  $(\diamond)$ .
  - If  $(\diamond)$  has solution  $\Phi$  strongly continuous  $\implies A = \Phi'(0)$  and  $\Phi(t) = \text{"exp}(tA)\text{"}$ .

# What we are going to show

## The example

we will construct  $X$  such that

$$\text{Iso}(X) = \{\pm \text{Id}\} \quad \text{but} \quad \text{Iso}(X^*) \supset \text{Iso}(\ell_2).$$

## The tools

- **Extremely non-complex Banach spaces:** spaces  $X$  such that  $\|\text{Id} + T^2\| = 1 + \|T^2\|$  for every  $T \in L(X)$ .
- **Koszmider type compact spaces:** topological compact spaces  $K$  such that  $C(K)$  has few operators.

## The talk is based on the papers



P. Koszmider, M. Martín, and J. Merí.  
Extremely non-complex  $C(K)$  spaces.  
*J. Math. Anal. Appl.* (2009).



P. Koszmider, M. Martín, and J. Merí.  
Isometries on extremely non-complex Banach spaces.  
*J. Inst. Math. Jussieu* (2011).



M. Martín  
The group of isometries of a Banach space and duality.  
*J. Funct. Anal.* (2008).

# Sketch of the talk

- 1 Introduction
- 2 Extremely non-complex Banach spaces:  
motivation and examples
- 3 Isometries on extremely non-complex spaces



## *Extremely non-complex Banach spaces: motivation and examples*

---

### Section 2

- 2 Extremely non-complex Banach spaces:  
motivation and examples
  - Complex structures
  - The first examples:  $C(K)$  spaces with few operators
  - More  $C(K)$ -type examples
  - Further examples

# Complex structures

## Definition

$X$  has **complex structure** if there is  $T \in L(X)$  such that  $T^2 = -\text{Id}$ .

## Some remarks

- This gives a structure of vector space over  $\mathbb{C}$ :

$$(\alpha + i\beta)x = \alpha x + \beta T(x) \quad (\alpha + i\beta \in \mathbb{C}, x \in X)$$

- Defining

$$\|x\| = \max\{\|e^{i\theta}x\| : \theta \in [0, 2\pi]\} \quad (x \in X)$$

one gets that  $(X, \|\cdot\|)$  is a complex Banach space.

- If  $T$  is an isometry, then the given norm of  $X$  is actually complex.
- Conversely, if  $X$  is a complex Banach space, then

$$T(x) = ix \quad (x \in X)$$

satisfies  $T^2 = -\text{Id}$  and  $T$  is an isometry.

## Complex structures II

### Some examples

- ① If  $\dim(X) < \infty$ ,  $X$  has complex structure iff  $\dim(X)$  is even.
- ② If  $X \simeq Z \oplus Z$  (in particular,  $X \simeq X^2$ ), then  $X$  has complex structure.
- ③ There are infinite-dimensional Banach spaces without complex structure:
  - **Dieudonné, 1952:** the James' space  $\mathcal{J}$  (since  $\mathcal{J}^{**} \equiv \mathcal{J} \oplus \mathbb{R}$ ).
  - **Szarek, 1986:** uniformly convex examples.
  - **Gowers-Maurey, 1993:** their H.I. space (no H.I. has complex structure) .

### Definition

$X$  is **extremely non-complex** if  $\text{dist}(T^2, -\text{Id})$  is the maximum possible, i.e.

$$\|\text{Id} + T^2\| = 1 + \|T^2\| \quad (T \in L(X))$$

Question (Gilles Godefroy, private communication, 2005)

Is there any  $X \neq \mathbb{R}$  such that  $\|\text{Id} + T^2\| = 1 + \|T^2\|$  for every  $T \in L(X)$ ?

## Examples: $C(K)$ spaces with few operators

### Theorem (Koszmider, 2004)

There are infinitely many different perfect compact spaces  $K$  such that all operators on  $C(K)$  are weak multipliers.

- They are called **weak Koszmider spaces**.

### Definition: weak multiplier

Let  $K$  be a compact space.  $T \in L(C(K))$  is a **weak multiplier** if

$$T^* = g\text{Id} + S$$

where  $g$  is a Borel function and  $S$  is weakly compact.

### Proposition

$K$  perfect,  $T \in L(C(K))$  weak multiplier  $\implies \|\text{Id} + T^2\| = 1 + \|T^2\|$

### Corollary

There are infinitely many non-isomorphic extremely non-complex spaces.

## More $C(K)$ -type examples

### More $C(K)$ type examples

There are perfect compact spaces  $K_1, K_2$  such that:

- $C(K_1)$  and  $C(K_2)$  are extremely non-complex,
- $C(K_1)$  contains a complemented copy of  $C(\Delta)$ .
- $C(K_2)$  contains a (1-complemented) isometric copy of  $\ell_\infty$ .

### Observation

- $C(K_1)$  and  $C(K_2)$  have operators which are not weak multipliers.
- They are decomposable spaces.

## Further examples

### Spaces $C_E(K\|L)$

$K$  compact,  $L \subset K$  closed nowhere dense,  $E \subset C(L)$ . Define

$$C_E(K\|L) := \{f \in C(K) : f|_L \in E\}.$$

### Observation

$$C_E(K\|L)^* \cong E^* \oplus_1 C_0(K\|L)^*$$

### Theorem

$K$  perfect weak Koszmider,  $L$  closed nowhere dense,  $E \subset C(L)$

$\implies C_E(K\|L)$  is extremely non-complex.

## *Isometries on extremely non-complex spaces*

---

### Section 3

- 3 Isometries on extremely non-complex spaces
  - Isometries on extremely non-complex spaces
  - Isometries on extremely non-complex  $C_E(K||L)$  spaces
  - The main example

# Isometries on extremely non-complex spaces

## Theorem

$X$  extremely non-complex.

- $T \in \text{Iso}(X) \implies T^2 = \text{Id}$ .
- $T_1, T_2 \in \text{Iso}(X) \implies T_1 T_2 = T_2 T_1$ .
- $T_1, T_2 \in \text{Iso}(X) \implies \|T_1 - T_2\| \in \{0, 2\}$ .
- $\Phi : \mathbb{R}_0^+ \longrightarrow \text{Iso}(X)$  one-parameter semigroup  $\implies \Phi(\mathbb{R}_0^+) = \{\text{Id}\}$ .

## Consequences

- $\text{Iso}(X)$  is a Boolean group with the composition.
- $\text{Iso}(X)$  identifies with the set  $\text{Unc}(X)$  of unconditional projections on  $X$ :

$$\begin{aligned}
 P \in \text{Unc}(X) &\iff P^2 = P, 2P - \text{Id} \in \text{Iso}(X) \\
 &\iff P = \frac{1}{2}(\text{Id} + T), T \in \text{Iso}(X), T^2 = \text{Id}.
 \end{aligned}$$



Extremely non-complex  $C_E(K\|L)$  spaces.

## Remember

$K$  perfect weak Koszmider,  $L$  closed nowhere dense,  $E \subset C(L)$   
 $\implies C_E(K\|L)$  is extremely non-complex and  $C_E(K\|L)^* \cong E^* \oplus_1 C_0(K\|L)^*$ .

## Proposition

$K$  perfect  $\implies \exists L \subset K$  closed nowhere dense with  $C[0,1] \subset C(L)$ .

## A good example

Take  $K$  perfect weak Koszmider,  $L \subset K$  closed nowhere dense with  
 $E = \ell_2 \subset C[0,1] \subset C(L)$ :

- $C_{\ell_2}(K\|L)$  has no non-trivial one-parameter semigroup of isometries.
- $C_{\ell_2}(K\|L)^* \cong \ell_2 \oplus_1 C_0(K\|L)^* \implies \text{Iso}(C_{\ell_2}(K\|L)^*) \supset \text{Iso}(\ell_2)$ .

But we are able to give a better result...

Isometries on extremely non-complex  $C_E(K\|L)$  spaces

## Theorem (Banach-Stone like)

$C_E(K\|L)$  extremely non-complex,  $T \in \text{Iso}(C_E(K\|L))$   
 $\implies$  exists  $\theta : K \setminus L \rightarrow \{-1, 1\}$  continuous such that

$$[T(f)](x) = \theta(x)f(x) \quad (x \in K \setminus L, f \in C_E(K\|L))$$

Consequence: cases  $E = C(L)$  and  $E = 0$ 

- $C(K)$  extremely non-complex,  $\varphi : K \rightarrow K$  homeomorphism  $\implies \varphi = \text{id}$
- $C_0(K \setminus L) \equiv C_0(K\|L)$  extremely non-complex,  $\varphi : K \setminus L \rightarrow K \setminus L$  homeomorphism  $\implies \varphi = \text{id}$

## Consequence: connected case

If  $K \setminus L$  is connected, then

$$\text{Iso}(C_E(K\|L)) = \{-\text{Id}, +\text{Id}\}$$

## The main example

Koszmider, 2004

$\exists \mathcal{K}$  weak Koszmider space such that  $\mathcal{K} \setminus F$  is connected if  $|F| < \infty$ .

Important observation on the construction above

There is  $\mathcal{L} \subset \mathcal{K}$  closed and nowhere dense, with

- $\mathcal{K} \setminus \mathcal{L}$  connected
- $C[0,1] \subseteq C(\mathcal{L})$

Consequence: the best example

Consider  $X = C_{\ell_2}(\mathcal{K}||\mathcal{L})$ . Then:

$$\text{Iso}(X) = \{-\text{Id}, +\text{Id}\} \quad \text{and} \quad \text{Iso}(X^*) \supset \text{Iso}(\ell_2)$$