



## Genetic study paves way for new neuropathic pain treatments

20/02/2018

Research news

A pioneering international study, carried out by the University of Granada, Harvard University, the University of California, Los Angeles (UCLA), the University of Florida, Laval University and the Jackson Laboratory has conducted an in-depth analysis of the molecular differences between the most common symptoms associated with neuropathic pain. The project may pave the way for the development of more effective painkillers for the treatment of this debilitating chronic condition, which afflicts approximately 500 million people throughout the world.



When an injury is incurred in the nervous system, this can give rise to a very specific condition known as neuropathic pain. The research study, conducted on mice, has examined the profile of the genetic changes that occur following these nerve injuries and aims to broaden our understanding of the mechanisms responsible for causing neuropathic pain.

When an individual suffers from this chronic condition, sensory stimuli such as temperature (especially cold sensations) and tactile sensations, which under normal circumstances do not generally cause pain, are interpreted by the nervous system as acutely painful. This abnormal response to stimuli is called allodynia.

Not all patients experience these painful symptoms with the same level of intensity, and some react more adversely than others to cold temperatures (cold allodynia) as opposed to tactile sensations (tactile allodynia), and vice versa.

This pioneering new study demonstrates that the causes of cold allodynia are exclusively neuronal in nature and that it is a direct response to the nerve injury, while tactile allodynia is the result of complex interactions between the immune system and the nervous system. Consequently, the diverse manifestations of <http://www.ugr.es/>

neuropathic pain have different cellular origins and therefore patients suffering from cold or tactile allodynia may be in need of different therapeutic strategies to treat their specific conditions.


According to Dr. Enrique J. Cobos del Moral, a Senior Lecturer at the Department of Pharmacology of the UGR and one of the main researchers behind the project, “taking an in-depth look at the molecular differences between the most common symptoms associated with neuropathic pain could lead to the development of more effective painkillers to alleviate this type of pain, designed to treat the specific symptoms experienced by each patient”.

This highly innovative research work, recently published in the prestigious journal **Cell Reports**, has been made possible thanks to the UGR Junior Doctoral Programme for New Lines of Research, supported by the Vice-Rectorate for Research and Knowledge Transfer of the UGR.

#### **Bibliographic reference:**

Cobos EJ, Nickerson C, Gao F, Chandran V, Bravo-Caparrós I, González-Cano R, Riva P, Andrews N, Latremoliere A, Seehus C, Perazzoli G, Nieto FR, Joller N, Painter M, Ma CHE, Omura T, Chesler EJ, Geschwind DH, Coppola G, Rangachari M, Woolf CJ, Costigan M. Mechanistic differences in neuropathic pain modalities revealed by correlating behavior with global expression profiling. *Cell Reports*, 22(5):1301-1312, 2018

Link to full article: [http://www.cell.com/cell-reports/fulltext/S2211-1247\(18\)30006-8](http://www.cell.com/cell-reports/fulltext/S2211-1247(18)30006-8)



Members of the UGR research group who developed the project. From left to right, Dr. Gloria Perazzoli, Dr. Francisco R. Nieto López, Inmaculada Bravo Caparrós and Dr. Enrique J. Cobos del Moral.

#### **Contact information:**

<http://www.ugr.es/>

Dr. Enrique J. Cobos del Moral

Department of Pharmacology of the University of Granada

Phone: +34 958 245 160

E-mail: @email

Compartir en