



## UGR scientists study the role of Mediterranean reservoirs as producers of greenhouse gases

21/07/2020

Research news

**Researchers from the Department of Ecology of the UGR have conducted the first-ever study of the carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) emissions of 12 reservoirs located in Granada, Jaén, and Córdoba in Andalusia**

A team of scientists from the Department of Ecology at the University of Granada (UGR) has conducted the first-ever study of the carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) emissions of 12 reservoirs located in Granada, Jaén, and Córdoba in Andalusia, Southern Spain. Their study demonstrated that reservoirs are important producers of greenhouse gases, contributing to the radiative forcing of aquatic ecosystems. They also found that the geology of the reservoir basin, the depth, and the human pressure on the basin all determine its climatic forcing (the potential to absorb heat in the atmosphere).

Human development has brought with it the construction of reservoirs to provide energy and supply water for human consumption and agricultural irrigation, especially in regions such as the Mediterranean. Spain is the European country with the highest number of built reservoirs. Such systems have a major social and environmental impact, and, recently, scientists have hypothesized that they act as major sources of greenhouse gases (GHGs). Studies have been conducted on GHG emissions from reservoirs in boreal and tropical regions, but, to date, there has been a paucity of data covering the Mediterranean biome, despite the proliferation of reservoirs in this area. Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide



(N<sub>2</sub>O) are the three most important GHGs of human origin. CO<sub>2</sub> is the most widely studied, while much less is known about CH<sub>4</sub> and N<sub>2</sub>O—yet they have a warming potential that is 34 and 298 times higher than CO<sub>2</sub> over a 100-year timescale, respectively.

In this study, which has been published in *Environmental Research Letters*, the UGR scientists observed that reservoirs located in calcareous areas emitted more CO<sub>2</sub> into the atmosphere than those located in siliceous areas that, at times, could even be CO<sub>2</sub> sinks. All the reservoirs emitted CH<sub>4</sub>, which was related to water temperature, reservoir depth, and eutrophication (that is, systems that become 'green' due to receiving excess nutrients).

The shallowest, 'greenest' systems with the highest water temperature emitted the most methane—a phenomenon that was particularly marked in the case of Cubillas reservoir in Granada. N<sub>2</sub>O emissions were very high in systems located in agricultural and urban areas that gave rise to large nitrogen inputs. The highest N<sub>2</sub>O emissions of all were obtained in the Iznájar reservoir in Córdoba. Watersheds located in areas with a forest cover of over 40% acted as N<sub>2</sub>O sinks.

The researchers combined the data on emissions of the three gases to obtain their radiative potential, based on their CO<sub>2</sub> equivalents. Overall, they found that Cubillas reservoir presented the highest values for radiative forcing per square metre, while, by total area of the reservoir, it was Iznájar that presented the highest radiative forcing. Furthermore, the forcing was much higher in summer than in winter. The authors highlight that the data from this research “helps us not only to determine the impact of the reservoirs that are already built but also to identify measures to help reduce emissions and even anticipate the impact of constructing future reservoirs. If we establish measures targeted at reducing the inputs of nutrients into aquatic systems, such as improving tertiary wastewater treatment and reducing the use of fertilizers, not only would this improve water quality but also reduce CH<sub>4</sub> and N<sub>2</sub>O emissions.”

The construction of reservoirs is currently witnessing a boom on a global level, especially in countries with emerging economies, where more than 3,000 large dams have been planned or are being built for the production of hydroelectric energy. It is for this reason that anticipating their impact on the climate is of vital importance. Wherever possible, the construction of reservoirs should take into account that siliceous and forested bedrocks and deep canyons can all help minimize GHG emissions.

This study was funded by the HERA Project (CGL2014-52362-R), the CRONOS Project (RTI2018-098849-B-I00), the Modeling Nature Scientific Unit (UCE.PP2017.03), the


Regional Ministry of Economy, Knowledge, Enterprise and Universities (CECEU) of Andalusia, and the European Regional Development Fund (ERDF, ref. SOMM17 SOMM17/6109/UGR). It forms part of the doctoral thesis of Elizabeth León-Palmero (FPU 014/02917), within the Doctoral Programme in Fundamental and Systems Biology at the University of Granada.

### **Bibliography:**


León-Palmero, E., Morales-Baquero, R., and Reche, I. (2020) 'Greenhouse gas fluxes from reservoirs determined by watershed lithology, morphometry, and anthropogenic pressure'. Environ. Res. Lett. 15 044012

Online: <https://iopscience.iop.org/article/10.1088/1748-9326/ab7467/meta>

### **Image captions:**



Eulogio Corral, laboratory technician from the Department of Ecology, University of Granada.



Elizabeth León-Palmero, researcher in the Department of Ecology, University of Granada, with the device that enables researchers to measure the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O flows simultaneously (PICARRO).

### **Media enquiries:**

Elizabeth León-Palmero

<http://www.ugr.es/>

Department of Ecology, University of Granada

Tel.: +34 958 248951

Email: [eleonpalmero@ugr.es](mailto:eleonpalmero@ugr.es)

Compartir en