

IMAGES OF CONTRACTIVE PROJECTIONS ON OPERATOR ALGEBRAS

LESLIE J. BUNCE AND ANTONIO M. PERALTA

ABSTRACT. It is shown that if P is a weak*-continuous contractive projection on a JBW*-triple M , then $P(M)$ is type I or semifinite, respectively, if M is of the corresponding type. We also show that $P(M)$ has no infinite spin part if M is a type I von Neumann algebra.

JW*-triples, that is, weak*-closed subspaces of $B(H)$ that are also closed under $x \mapsto xx^*x$, arise as images of contractive (i.e. norm one) projections on von Neumann algebras. Their generalisations, JBW*-triples, are those complex Banach dual spaces whose open unit ball is a bounded symmetric domain. The holomorphy of such spaces induces a ternary Jordan algebraic structure determined by a certain “triple product” $\{a, b, c\}$ [16]. If $P : M \rightarrow M$ is a weak*-continuous contractive projection on a JBW*-triple M then $P(M)$ is a JBW*-triple with a triple product given by $\{a, b, c\}_P := P\{a, b, c\}$ by [17], [19], and by [9, 10] if M is a JW*-triple. The interesting special cases that occur when P is positive unital acting on von Neumann algebra or a JBW*-algebra were studied earlier in [4], [7] and [18].

Suppose $P : M \rightarrow M$ is a weak*-continuous contractive projection on a JBW*-triple M . In this paper we study the stability of $P(M)$ with respect to the type theory of [13, 14, 15]. We show that if M is type I or semifinite, respectively, then $P(M)$ is of the corresponding type. This extends the classical results of [22] when M is a von Neumann algebra and $P(M)$ is a subalgebra. We remark that in general $P(M)$ is not a subtriple of M . Using recent results on properties of the predual of a type I von Neumann algebra we deduce that $P(M)$ cannot be isometric to an infinite dimensional spin factor whenever M is a type I von Neumann algebra.

The first section of this paper contains preliminary results on JBW*-algebras. This is continued in §2 where we study the fixed point JW*-algebra, W^α , of an involution α on a von Neumann algebra W . A principal aim here is to show that a faithful weak*-continuous contractive projection from W^α onto a continuous JW*-subalgebra induces a weak*-continuous contractive projection from W onto a continuous von Neumann subalgebra. This allows us to apply [22] to obtain our main results in §4. The formulation

2000 *Mathematics Subject Classification*. Primary 46B04, 46B20, 46L05, and 46L10.

Second author partially supported by D.G.I.C.Y.T. project no. PB 98-1371, and Junta de Andalucía grant FQM 0199 .

of type theory of JBW*-triples contained in §3 is extracted from [13, 14, 15] and is included for completeness.

For later reference we shall recall some of the fundamentals of JBW*-triples used in this paper. A JBW*-triple can be realised [16] as a complex Banach space M with predual M_* and continuous ternary triple product $(a, b, c) \mapsto \{a, b, c\}$ that is conjugate linear in b and symmetric bilinear in a, c such that $\|\{a, a, a\}\| = \|a\|^3$ and such that the operator $x \mapsto \{a, a, x\}$, denoted by $D(a, a)$, is hermitian with non-negative spectrum and satisfies

$$D(a, a)(\{x, y, z\}) = \{D(a, a)x, y, z\} - \{x, D(a, a)y, z\} + \{x, y, D(a, a)z\}.$$

The predual is unique and the triple product is separately weak*-continuous [2], [13]. The surjective linear isometries between JBW*-triples are the triple product preserving bijections (triple isomorphisms) [16]. A von Neumann algebra is a JBW*-triple with triple product $\{a, b, c\} = \frac{1}{2}(ab^*c + cb^*a)$. The weak*-closed subtriples of von Neumann algebras are the JW*-triples. By [14, 15] most JBW*-triples are of this form. See §3 for further details.

An element u in a JBW*-triple M satisfying $\{u, u, u\} = u$ is called a *tripotent*, when M is a JW*-triple these are precisely the partial isometries of M . Associated with a tripotent u are the mutually orthogonal *Peirce* projections $P_2(u)$, $P_1(u)$, and $P_0(u)$. We have,

$$P_2(u)(x) = \{u, \{u, x, u\}, u\} \quad \text{for all } x,$$

$$P_1(u) = 2(D(u, u) - P_2(u)) \quad \text{and} \quad P_2(u) + P_1(u) + P_0(u) = i$$

(where i is the identity map). A tripotent u of M is said to be *complete* (or maximal) if $P_0(u) = 0$, to be *unitary* if $P_2(u) = i$ and to be *minimal* if $P_2(u)(M) = \mathbb{C}u$. We recall (see [5, Corollary 4.8], for example) that the complete tripotents of M are the extreme points of the closed unit ball of M . A crucial simplifying property of JBW*-triples is that for a tripotent u of M the Peirce-2-subspace $P_2(u)(M)$ is a JBW*-algebra with product $a \circ b = \{a, u, b\}$ and involution $a^* = \{u, a, u\}$. For further properties of JBW*-triples we refer to the papers [9], [5, 6], [13, 14, 15] and [16], and the book [24]. Since JBW*-algebras are just the complexifications of JW-algebras we refer to [12] for their theory.

1. POSITIVE UNITAL PROJECTIONS ON JBW*-ALGEBRAS

Let M be a JBW*-algebra. Writing

$$[a, b, c] := (a \circ b) \circ c + (c \circ b) \circ a - (a \circ c) \circ b,$$

M is a JBW*-triple with triple product given by $\{a, b, c\} := [a, b^*, c]$. The Peirce-2 projection, $P_2(e)$, associated with a projection e of M satisfies $P_2(e)(x) = [e, x, e]$ for all x in M .

Elements a and b of M are said to *operator commute* in M if $(a \circ x) \circ b = a \circ (x \circ b)$ for all x in M . Self-adjoint elements a and b in M generate a JBW*-subalgebra that can be realised as a JW*-subalgebra of some $B(H)$ [25] and, in this realisation, a and b commute in the usual

sense if they operator commute in M [23, Proposition 1]. By the same references, self-adjoint elements a and b of M operator commute if and only if $a^2 \circ b = [a, b, a] (= \{a, b, a\})$. If N is a JBW^* -subalgebra of M we use $M \cap N'$ to denote the set of elements in M that operator commute with every element of N . (This corresponds to the usual notation when M is a von Neumann algebra). The *centre* of M is $M \cap M'$ which we also denote by $Z(M)$.

Let P be a unital (i.e. $P(1) = 1$) weak*-continuous contractive projection on a JBW^* -algebra M . Then P is positive and therefore is invariant on the self-adjoint part. Such projections were studied in [7] and [18]. Suppose now that $P(M)$ is a JBW^* -subalgebra N of M . Then, by [7, Lemma 1.5], [18, Lemma 1.5] we have $P(a \circ x) = a \circ P(x)$ for all $a \in N$ and $x \in M$. Further, if e denotes the support projection of P in M (i.e. the least projection in M sent to 1 by P) then $P = PP_2(e)$ and, by a slight extension of [7, Lemma 1.2(2)], $e \in M \cap N'$. Moreover, if $x \geq 0$ and $P(x) = 0$ then $P_2(e)(x) = 0$. If $e = 1$, P is said to be faithful.

Lemma 1.1. *Let $P : M \rightarrow M$ be a weak*-continuous unital contractive projection from a JBW^* -algebra M onto a JBW^* -subalgebra N . Let e be the support projection of P . Then $P_2(e)P$ is a faithful weak*-continuous unital projection from $P_2(e)(M)$ onto $N \circ e$. Moreover, N is isomorphic to $N \circ e$.*

Proof. Suppose $x \in P_2(e)(M)$ such that $x \geq 0$ and $P_2(e)P(x) = 0$. Then $P(x) = PP_2(e)P(x) = 0$ so that $x = P_2(e)(x) = 0$. Together with the above remarks this proves the first statement.

Since $e \in M \cap N'$ multiplication by e induces a (Jordan) homomorphism, π , from N onto $N \circ e$. Let a in N such that $a \geq 0$ and $a \circ e = 0$, then $a = P(a \circ e) = 0$. It follows that π is injective. \square

Lemma 1.2. *Let $P : M \rightarrow M$ be a weak*-continuous unital contractive projection from a JBW^* -algebra onto a JBW^* -subalgebra N . Let e be any non-zero projection in $M \cap N'$. Suppose that P is faithful. Then there exists a faithful weak*-continuous unital contractive projection from $P_2(e)(M)$ onto $N \circ e$.*

Proof. For each self-adjoint $a \in N$ we have

$$a^2 \circ P(e) = P(a^2 \circ e) = P(\{a, e, a\}) = \{a, P(e), a\}$$

and so, by the previous remark, $P(e) \in Z(N)$. Therefore the range projection $r(P(e)) \in Z(N)$. Denote $r(P(e))$ by h . The ideal of N , $N \circ P(e) = P(N \circ e)$, is weak*-closed and so equals $N \circ h$. It follows that $P(e)$ is invertible in $N \circ h$ with inverse b , say, in $Z(N) \circ h$. Define $Q : P_2(e)(M) \rightarrow P_2(e)(M)$ by $Q(x) = (P(x) \circ b) \circ e$. Let $a \in N$ where $a \geq 0$. By operator commutivity we have $(a \circ (1 - h)) \circ e \geq 0$ and

$$P((a \circ (1 - h)) \circ e) = (a \circ (1 - h)) \circ P(e) = a \circ ((1 - h) \circ P(e)) = 0.$$

Since P is faithful, $a \circ e = (a \circ h) \circ e$, and so

$$Q(a \circ e) = (P(a \circ e) \circ b) \circ e = ((a \circ P(e)) \circ b) \circ e = (a \circ h) \circ e = a \circ e,$$

implying that Q is a unital projection onto $N \circ e$. To see that Q is faithful let $x \in P_2(e)(M)$ such that $x \geq 0$ and $(P(x) \circ b) \circ e = 0$. By the above,

$$P(x) \circ e = (P(x) \circ h) \circ e = ((P(x) \circ b) \circ e) \circ P(e) = 0.$$

Therefore, $P(x) \circ P(e) = P(P(x) \circ e) = 0$. But $P(x) \leq \|x\|P(e)$. Hence, $P(x) = 0$ and so $x = 0$ because P is faithful. \square

2. INVOLUTORY * ANTIAUTOMORPHISMS

Following [21] by an involution α on a von Neumann algebra we shall mean an involutory * antiautomorphism on the algebra. Let α be an involution on a von Neumann algebra W . We shall write $R(W) := \{x \in W : \alpha(x) = x^*\}$ and $W^\alpha := \{x \in W : \alpha(x) = x\}$ (The latter notation is different from that used in [21], where it stands for the hermitian part). Then $R(W)$ is a weak*-closed real *-subalgebra of W with $R(W) \cap iR(W) = \{0\}$ and $W = R(W) + iR(W)$. We have $W^\alpha = R(W)_{sa} + iR(W)_{sa}$ and, for $a, b \in R(W)$, we have $\alpha(a + ib) = a^* + ib^*$.

Lemma 2.1. *Let α be an involution on a von Neumann algebra W and suppose e is a central projection in W such that $e + \alpha(e) = 1$. Then $eW^\alpha = eW$ and W^α is (Jordan) isomorphic to eW via $x \mapsto ex$.*

Proof. For each x in W , $ex + (1 - e)\alpha(x) \in W^\alpha$ and every element of W^α is of this form. Thus $eW^\alpha = eW$ and W^α is isomorphic to eW in the way stated. \square

Lemma 2.2. *Let α be an involution on a von Neumann algebra W and suppose that e is a projection in W with $e + \alpha(e) = 1$. Then we have the following.*

- (i) *There is a faithful weak*-continuous unital contractive projection, $P : W^\alpha \rightarrow W^\alpha$, such that $P(W^\alpha)$ is a JW^* -subalgebra isomorphic to eWe (and to $(1 - e)W(1 - e)$).*
- (ii) *If W^α generates W as a von Neumann algebra and $eW^\alpha\alpha(e) = 0$, then $e \in Z(W)$.*

Proof. (i) Let V denote the von Neumann algebra $eWe + (1 - e)W(1 - e)$. Define $P : W \rightarrow W$ by $P(x) := exe + (1 - e)x(1 - e)$. Then $P(W) = V = \alpha(V)$. If s denotes the symmetry $2e - 1$ we see that $P(x) = \frac{1}{2}(x + sxs)$. Since $\alpha(s) = -s$, we have $\alpha P = P\alpha$ from which we deduce that $P(W^\alpha) = V^\alpha$. Since e lies in the centre of V , Lemma 2.1 implies that V^α is isomorphic to $eV = eWe$. It is clear that P satisfies (i).

(ii) Suppose $eW^\alpha\alpha(e) = 0$. Then for $x \in W^\alpha$ we have

$$x = exe + (1 - e)x(1 - e)$$

so that $ex = exe$. Passing to the self-adjoint part we see that e commutes with all elements of W^α and so lies in the centre of W if W is the von Neumann algebra generated by W^α . \square

Lemma 2.3. *Let α be an involution in a non-abelian von Neumann algebra W . Then there is a non-zero projection e in W with $e\alpha(e) = 0$.*

Proof. We have $R(W)_{sa} \neq R(W)$ otherwise α is the identity map on W and therefore W is abelian. Choose a in $R(W)$ such that $a \neq a^*$ and let $a - a^* = b$. Let V denote the von Neumann subalgebra of W generated by b . We have that V is abelian, that $\alpha(b) = -b$ and $\alpha(V) = V$. Since α is not the identity map on V , by [12, 7.3.4] there is a non-zero projection $e \in V$ such that $e\alpha(e) = 0$. \square

Proposition 2.4. *Let α be an involution on a von Neumann algebra W and suppose that W^α has no type I part. Then there is a projection e in W and a faithful weak*-continuous unital contractive projection from W^α onto a JW*-subalgebra M such that $e \in W \cap M'$ and Me is a W^* -algebra isomorphic to M .*

Moreover, if W^α is of type II_1 , II_∞ or III , respectively, then M is of the corresponding type.

Proof. Let (e_i) be a family of projections in W maximal subject to the condition that $(e_i + \alpha(e_i))$ is a mutually orthogonal family of projections. Put $e = \sum_i e_i$. Then $e\alpha(e) = 0$. Let $f = 1 - e - \alpha(e)$. Then $\alpha(fWf) = fWf$ and it follows from Lemma 2.3, and maximality, that fWf is abelian and hence that $fW^\alpha f$ is abelian. By assumption, we must have $f = 0$. Lemma 2.2(i) now gives the first statement. Since W^α generates W , by [11, Theorem 2.8], the second statement follows from [1, Theorem 8] together with Lemma 2.2(i). \square

Proposition 2.5. *Let α be an involution on a von Neumann algebra W , and let M denote W^α . Suppose there is a faithful weak*-continuous unital contractive projection, P , from M onto a JW*-subalgebra N . If N is continuous (respectively, type III) then there is a weak*-continuous contractive projection from W onto a continuous (respectively type III) W^* -subalgebra.*

Proof. Let V the von Neumann subalgebra of W generated by N and let R be the weak*-closed real *-subalgebra of W generated by V_{sa} . We have $\alpha(V) = V$ since α fixes each element of N , and $R \cap iR = \{0\}$ since $R \subset R(W)$. Suppose N is continuous (respectively, type III). Then $N_{sa} = R_{sa}$, using [12, 7.3.3], so that $V = R + iR$, by [20, Theorem 2.4]. Hence, $V^\alpha = R_{sa} + iR_{sa} = N$. By Proposition 2.4 there exists a faithful weak*-continuous unital contractive projection, $Q : N \rightarrow N$, onto a continuous (respectively, type III) JW*-subalgebra K together with a projection $e \in W \cap K'$ such that Ke is a W^* -algebra isomorphic to K . If E denotes

the (faithful) canonical projection $\frac{1}{2}(i + \alpha) : W \rightarrow M$, then the proof is completed by application of Lemma 1.2 to the projection $QPE : W \rightarrow K$. \square

We recall ([21]) that an involution α is said to be a *central* involution if it fixes every element in $Z(W)$.

Lemma 2.6. *Let α be a central involution on a continuous von Neumann algebra W . Let u be a partial isometry of W^α such that $(1 - uu^*)W^\alpha(1 - u^*u) = 0$. Then $u^*u = uu^* = 1$.*

Proof. Let e denote $1 - uu^*$. Then $\alpha(e) = 1 - u^*u$. Put $p = e + \alpha(e)$. Then α is a central involution on pWp . By [11, Theorem 2.8] on [21, Proposition 3.2] $(pWp)^\alpha (= pW^\alpha p)$ generates pWp . Hence by Lemma 2.2(ii), $e \in Z(pWp) = Z(W)p$ so that $\alpha(e) = e$, whence the result. \square

3. TYPES OF JBW*-TRIPLES

The aim of this short section, which contains no new results, is to collate existing theory into a form easy to use subsequently.

Cartan factors. Of the six kinds of Cartan factors (up to linear isometry), three are of the form $pB(H)$, $\{x \in B(H) : x = jx^*j\}$ and $\{x \in B(H) : x = -jx^*j\}$, where H is a complex Hilbert space, p is a projection in $B(H)$ and $j : H \rightarrow H$ is a conjugation. These are referred to as *rectangular*, *hermitian* and *symplectic* Cartan factors, respectively. Hermitian factors are type I JW*-algebra factors and, if H is even or infinite dimensional, symplectic factors are linearly isometric to type I JW*-algebra factors. *Spin* factors (complexifications of real spin factors) comprise a fourth kind. The remaining two *exceptional* Cartan factors can be realized as the 3×3 hermitian matrices and the 1×2 matrices, respectively, over the complex Cayley numbers.

Type I JBW-triples.* In view of [13, 4.14] a JBW*-triple M is said to be *type I* if there is a complete tripotent u of M such that $P_2(u)(M)$ is a type I JBW*-algebra. By the type I classification theorem [14, 1.7] the type I JBW*-triples are precisely the ℓ_∞ -sums of JBW*-triples of the form

- (i) : $A\overline{\otimes}C$, where A is an abelian von Neumann algebra and C is a Cartan factor realised as a JW*-subtriple of some $B(H)$, the bar denoting the weak*-closure in the usual von Neumann tensor product $A\overline{\otimes}B(H)$, and
- (ii) : $A \otimes C$ (algebraic tensor product) where A is as before and C is an exceptional Cartan factor.

(Of course, $A\overline{\otimes}C = A \otimes C$ whenever C is a finite dimensional non-exceptional Cartan factor.)

Let e be a tripotent in a type I JBW*-triple M . A known consequence of the type I classification theorem is that $P_2(e)(M)$ is type I. We include an argument for completeness and want of a precise reference.

We may suppose that M is of the form (i) or (ii), above. In the latter case it is clear that $P_2(e)(M)$ is type I since every subfactor of it must have rank less than 4. Thus we may assume that we are in the case (i) and, consequently, that we are working in $A\overline{\otimes}B(H)$.

Let u be a nonzero (we assume $e \neq 0$) in a weak*-closed ideal J of $P_2(e)(M)$.

Since $\{u, (A\otimes B(H)), u\} = (A\otimes 1)\{u, (1\otimes B(H)), u\}$ and $B(H)$ is the weak*-closed linear span of its minimal tripotents, $\{u, (1\otimes v), u\} \neq 0$ for some minimal tripotent v . We have $\{(1\otimes v), M, (1\otimes v)\} = A\otimes v$ so that with $x = \{u, (1\otimes v), u\} \in P_2(u)(M)$ we have $\{x, M, x\} \subset (A\otimes 1)x$. Since $A\otimes 1$ commutes elementwise with x , $(A\otimes 1)x$ generates an abelian subtriple in the sense of [13, 1.4]. But, as follows from [5, Lemma 3.1], the weak*-closure of $\{x, M, x\}$ equals $P_2(w)(M)$, for some tripotent w , and so is abelian. Since $w \in J$, $P_2(e)(M)$ is type I, by [13, 4.14 (2) \Rightarrow (1)].

Continuous JBW-triples.* A JBW*-triple M is said to be *continuous* if it has no type I ℓ_∞ -summand. In which case, up to isometry, M is a JW*-triple with unique decomposition, $M = W^\alpha \oplus pV$, where W and V are continuous von Neumann algebras, p is a projection in V and α is a *central* involution on W [15, 2.1 and 4.8]. It is implicit in [15] that every complete tripotent of W^α is a unitary tripotent. An alternative proof of this fact is provided by Lemma 2.6. Thus, by [15, 5.1-5.7], for every complete tripotent u in M , $P_2(u)(M)$ is isometric to $W^\alpha \oplus pWp$. We define M to be of type II_1 , II_∞ or III , respectively if both W and pWp are of the corresponding type. M is said to be *semifinite* if it has no type III ℓ_∞ -summand.

Lemma 3.1, below, summarizes the above. The second statement is a consequence of the fact that every tripotent in a JBW*-triple M is a projection in $P_2(u)(M)$ for some complete tripotent u [13, 3.12].

Lemma 3.1. *A JBW*-triple M is of type I, II_1 , II_∞ , III or is semifinite, respectively if and only if $P_2(u)(M)$ is of the corresponding type for some, and hence every, complete tripotent u of M . If M is of type I, II_1 , III or is semifinite, respectively, then so is $P_2(u)(M)$ for every tripotent u of M .*

We shall say that a JBW*-triple has no *infinite spin part* if it has no ℓ_∞ -summands of the form $A\overline{\otimes}C$ where A is an abelian von Neumann algebra and C is an infinite dimensional spin factor.

4. CONTRACTIVE PROJECTIONS ON JBW*-TRIPLES

By [17] and [19] the image of a weak*-continuous contractive projection, $P : M \rightarrow M$, on a JBW*-triple M is again a JBW*-triple with triple product $\{x, y, z\}_P := P(\{x, y, z\})$ for x, y, z in $P(M)$ and

$$P\{P(x), y, P(z)\} = P\{P(x), P(y), P(z)\}$$

for all x, y, z in M . The image, $P(M)$, need not be a JBW*-subtriple of M . However, as is made explicit in [6, Lemma 5.3] and its proof, we do have the following.

Lemma 4.1. [6, Lemma 5.3] *If $P : M \rightarrow M$ is a weak*-continuous contractive projection on a JBW*-triple M , there exists a JBW*-subtriple C of M such that C is linearly isometric to $P(M)$ and such that C is the image of a weak*-continuous projection on M .*

We are now in a position to prove our first main result. We freely use Lemma 3.1 throughout.

Theorem 4.2. *Let $P : M \rightarrow M$ be a weak*-continuous contractive projection on a JBW*-triple M . If M is of type I (respectively, semifinite) then $P(M)$ is type I (respectively, semifinite).*

Proof. Let M be type I (respectively, semifinite). By Lemma 4.1 we may suppose $P(M)$ to be a JBW*-subtriple, N , of M . Let u be a complete tripotent of N . By the above formula, P restricts to a unital projection from $P_2(u)(M)$ to $P_2(u)(N)$.

By this fact, together with Lemma 1.1, we may suppose P to be faithful, M to be a JBW*-algebra and N to be a JBW*-subalgebra.

Let $M \circ z$ be the type I finite part of M , where z is a central projection of M . Then $N \circ z$ is type I finite, being a subalgebra of $M \circ z$, and it remains only to show that $N \circ (1 - z)$ is type I (respectively, semifinite). Since, by Lemma 1.2, $N \circ (1 - z)$ is the image of some faithful weak*-continuous unital contractive projection on $M \circ (1 - z)$, it can be supposed that $z = 0$. In which case, by [12, 7.2.7 and 7.3.3], we may suppose that $M = W^\alpha$, where α is an involution on a von Neumann algebra W . Since W^α generates W [11, Theorem 2.8], W is type I (respectively, semifinite) by [12, 7.4.2] and [1, Theorem 8].

In order to obtain a contradiction, suppose now that N has a non-zero continuous (respectively, type III) part, $N \circ e$, where e is a central projection of N . Now, α is an involution on eWe with $(eWe)^\alpha = eMe$. Applying Proposition 2.5 to $P : eMe \rightarrow N \circ e$, which is surjective, we obtain a weak*-continuous projection from the type I (respectively, semifinite) W*-algebra eWe onto a continuous (respectively, type III) W*-subalgebra. This contradicts [22, Theorem 3 (respectively, Theorem 4)] and so completes the proof. \square

In order to prove a refinement of part of Theorem 4.2, we first recall a Banach space property introduced in [8].

Definition. A Banach space E is said to have the DP1 if whenever a sequence $x_n \rightarrow x$ weakly in E with $\|x_n\| = \|x\| = 1$ for all n , and (ρ_n) is a weakly null sequence in E^* , then $\rho_n(x_n) \rightarrow 0$.

We write M_* for the predual of a JBW*-triple M and we note that if $P : M \rightarrow M$ is a weak*-continuous contractive projection then the dual projection restricts to a contractive projection on M_* and that $P(M)_*$ is linearly isometric to $P^*(M_*)$ via $\tau \mapsto \tau \circ P$. It follows that if M_* has the DP1 then so does $P(M)_*$.

Recently, the authors characterised the von Neumann algebras whose predual has the DP1.

Lemma 4.3. [3, Theorem 6] *A von Neumann algebra is type I if and only if its predual has the DP1.*

For properties of (real) spin factors used in the next proof, see [12, §6].

Lemma 4.4. *Let C be an infinite dimensional spin factor. Then C_* does not have the DP1.*

Proof. The argument is similar to that in [3, Proposition 5]. Let τ denote the tracial state of C and let R be the real Banach space generated by the non-trivial symmetries in C . Then R is isometric to an infinite dimensional real Hilbert space and $\tau(R) = \{0\}$. Let (s_n) be an infinite orthogonal sequence in the Hilbert space R . Then $(s_n) \rightarrow 0$ weakly in R and hence in C . Moreover, each s_n is a non-trivial symmetry. For each n , let e_n denote the projection $\frac{1}{2}(1 + s_n)$ and let τ_n denote the normal state $2\tau(e_n \cdot e_n)$. For all n , $e_n s_n e_n = e_n$ so that $\tau(s_n) = 1$. However, $\tau_n \rightarrow \tau$ weakly in C_* , since $\tau_n(x) = 2\tau_n(e_n \circ x)$, for all x and n . Therefore, C_* does not have the DP1. \square

One immediate consequence of Lemma 4.4 is that if A is an abelian von Neumann algebra and C is an infinite dimensional spin factor then (in the notation of §3) $(A \overline{\otimes} C)_*$ cannot have the DP1 because of the canonical (weak*-continuous) contractive projection $A \overline{\otimes} C \rightarrow C$.

Theorem 4.5. *Let M be a JBW*-triple. Then M_* has the DP1 if and only if M is type I without infinite spin part.*

Proof. Suppose M_* has the DP1. Then the predual of every ℓ_∞ -summand of M has the DP1. Thus by Proposition 2.4 and Lemma 4.3, M cannot have a non-zero ℓ_∞ -summand of the form W^α where α is an involution on a continuous von Neumann algebra W , nor of the form pV where p is a non-zero projection in a continuous von Neumann algebra V . (In the latter case because of the natural projection $pV \rightarrow pVp$). Therefore, M is type I and, by the remark prior to the statement of the theorem, has no infinite spin part.

On the other hand, consider an abelian von Neumann algebra A and a Cartan factor C . If C is finite dimensional then $A \otimes C$ has the Dunford-Pettis property because A does, and so $(A \otimes C)_*$ has the Dunford-Pettis property and therefore it has the DP1. Suppose C is infinite dimensional.

If C is (rectangular) of the form $pB(H)$ for a projection $p \in B(H)$, then $A\overline{\otimes}C = (1 \otimes p)A\overline{\otimes}C$ and is clearly the image of a weak*-continuous projection on $A\overline{\otimes}B(H)$, implying that $(A\overline{\otimes}C)_*$ has the DP1, by Lemma 4.3. If C is hermitian or symplectic then $A\overline{\otimes}C$ can be realised as W^α where α is an involution on a type I von Neumann algebra W , by [12, 7.3.3]. Since W^α is the image of the weak*-continuous contractive projection $\frac{1}{2}(i + \alpha)$ on W , Lemma 4.3 again gives that $(A\overline{\otimes}C)_*$ has the DP1. Thus, if M is type I with no infinite spin part, M_* has the DP1 by [8, 1.10] together with [14, 1.7]. \square

This leads to the following refinement of Theorem 4.2. The proof is immediate from Theorem 4.5.

Theorem 4.6. *Let $P : M \rightarrow M$ be a weak*-continuous contractive projection on a JBW*-triple M where M is type I with no infinite spin part. Then $P(M)$ is type I with no infinite spin part.*

For every spin factor C acting on a complex Hilbert space H there is a positive unital projection from $B(H)$ onto C [7, Lemma 2.3]. Since a von Neumann algebra never has infinite spin part, Theorem 4.6 gives:

Corollary 4.7. *There is no weak*-continuous contractive projection from a type I von Neumann algebra onto an infinite dimensional spin factor.*

REFERENCES

- [1] L. Ajupov, Extensions of traces and type criterions for Jordan algebras of self adjoint operators, *Math. Z.* **181** (1982), 253-268.
- [2] J. T. Barton and R. Timoney, Weak* continuity of Jordan triple product and its applications, *Math. Scand.* **59** (1986), 177-191.
- [3] L. J. Bunce and A. M. Peralta, The alternative Dunford-Pettis property in C*-algebras and von Neumann preduals, preprint, 2001.
- [4] M. Choi and E. Effros, Injectivity in operator spaces, *J. Funct. Analysis* **24** (1977), 156-209.
- [5] C. M. Edwards and G. T. Rüttiman, On the facial structure of the unit balls in a JBW*-triple and its predual, *J. London Math. Soc.* **38** (1988), 317-332.
- [6] C. M. Edwards and G. T. Rüttiman, Structural projections on JBW*-triples, *J. London Math. Soc.* **53** (1996), 354-368.
- [7] E. Effros and E. Størmer, Positive projections and Jordan structure in operator algebras, *Math. Scand.* (1983), 279-311.
- [8] W. Freedman, An alternative Dunford-Pettis property, *Studia Math.* **125** (1997), 143-159.
- [9] Y. Friedman and B. Russo, Conditional expectations without order, *Pac. J. Math.* **115** (1984), 351-360.
- [10] Y. Friedman and B. Russo, Solution of the contractive projection problem, *J. Funct. Analysis* **60** (1986), 56-79.
- [11] J. Gasmyr, Involutory *-antiautomorphisms of von Neumann algebras and C*-algebras, *Math. Scand.* **67** (1990), 87-96.
- [12] Hanche-Olsen, H. and Størmer, E., *Jordan operator algebras*, Pitman, London, 1984.

- [13] G. Horn, Characterization of the predual and ideal structure of a JBW^* -triple, *Math. Scand.* **61** (1987), 117-133.
- [14] G. Horn, Classification of JBW^* -Triples of type I, *Math. Z.* **196** (1987), 271-291.
- [15] G. Horn and E. Neher, Classification of continuous JBW^* -Triples, *Trans. Amer. Math. Soc.* **306** (1988), 553-578.
- [16] W. Kaup, A Riemann mapping theorem for bounded symmetric domains in complex Banach spaces, *Math. Z.* **183** (1983), 503-529.
- [17] W. Kaup, Contractive projections on Jordan C^* -algebras and generalizations, *Math. Scand.* **54** (1984), 95-100.
- [18] A. G. Robertson and M. A. Youngson, Positive projections with contractive complements on Jordan algebras, *J. London Math. Soc.* **26** (1982), 132-142.
- [19] L. L. Stacho, A projection principle concerning biholomorphic automorphisms, *Acta Sci. Math.* **44** (1982), 99-124.
- [20] E. Størmer, Irreducible Jordan algebras of self-adjoint operators, *Trans. Amer. Math. Soc.* **130** (1966), 153-166.
- [21] E. Størmer, Conjugacy of involutive antiautomorphisms of von Neumann algebras, *J. Funct. Analysis* **66** (1986), 54-66.
- [22] J. Tomiyama, On the projection of norm one in W^* -algebras III, *Tohoku Math. J.* **11** (1959), 125-129.
- [23] D. Topping, *Jordan algebras of self-adjoint operators*, Mem. Amer. Math. Soc. **53**, 1965.
- [24] H. Upmeyer, *Symmetric Banach manifolds and Jordan C^* -algebras*, Math. Studies 104, North Holland, 1985.
- [25] J. D. M. Wright, Jordan C^* -algebras, *Mich. Math. J.* **24** (1977), 291-302.

UNIVERSITY OF READING, READING RG6 2AX, GREAT BRITAIN.

E-mail address: L.J.Bunce@reading.ac.uk

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA 18071 GRANADA, SPAIN.

E-mail address: aperalta@goliat.ugr.es